

Harmonic functions and instanton moduli spaces on the multi-Taub–NUT space

Gábor Etesi

*Department of Geometry, Mathematical Institute, Faculty of Science,
Budapest University of Technology and Economics,
Egry J. u. 1, H ép., H-1111 Budapest, Hungary*^{*}

Szilárd Szabó

*Department of Geometry, Mathematical Institute, Faculty of Science,
Budapest University of Technology and Economics,
Egry J. u. 1, H ép., H-1111 Budapest, Hungary*[†]

and

*Rényi Institute of Mathematics, Hungarian Academy of Science,
Reáltanoda u. 13-15., H-1053 Budapest, Hungary*[‡]

September 2, 2008

Abstract

Explicit construction of the basic $SU(2)$ anti-instantons over the multi-Taub–NUT geometry via the classical conformal rescaling method is exhibited. These anti-instantons satisfy the so-called weak holonomy condition at infinity with respect to the trivial flat connection and decay rapidly. The resulting unitary energy anti-instantons have trivial holonomy at infinity.

We also fully describe their unframed moduli space and find that it is a five dimensional space admitting a singular disk-fibration over $\mathbb{R}^3$.

On the way, we work out in detail the twistor space of the multi-Taub–NUT geometry together with its real structure and transform our anti-instantons into holomorphic vector bundles over the twistor space. In this picture we are able to demonstrate that our construction is complete in the sense that we have constructed a full connected component of the moduli space of solutions of the above type.

We also prove that anti-instantons with arbitrary high integer energy exist on the multi-Taub–NUT space.

AMS Classification: Primary: 53C07; Secondary: 14F05, 53C28, 58J99

Keywords: *Multi-Taub–NUT space; Yang–Mills instantons; Harmonic functions; Twistors*

^{*}etesi@math.bme.hu

[†]szabosz@math.bme.hu

[‡]szabosz@renyi.hu

1 Introduction

The aim of this paper is to construct the most relevant anti-instanton moduli space over the multi-Taub–NUT geometry by elementary means.

An important class of non-compact but complete four dimensional geometries is the collection of the so-called *asymptotically locally flat* (ALF) spaces including several mathematically as well as physically important examples. The flat $\mathbb{R}^3 \times S^1$ plays a role in finite temperature Yang–Mills theories, the Euclidean Schwarzschild space [20] deals with quantum gravity and Hawking radiation. If the metric is additionally hyper-Kähler then the space is also called an *ALF gravitational instanton* (in the narrow sense). The flat $\mathbb{R}^3 \times S^1$ is a straightforward example; non-trivial ones for this restricted class are provided by the multi-Taub–NUT (or A_k ALF or ALF Gibbons–Hawking) spaces [15] which carry supersymmetric solutions of string theory and supergravity models, the Atiyah–Hitchin manifold (and its universal double cover) [3] describing the 2-monopole moduli space over $\mathbb{R}^3$ and last but not least the recently constructed D_k ALF spaces [9].

The ALF asymptotics is a natural generalization of the well-known ALE (asymptotically locally Euclidean) one including the multi-Eguchi–Hanson geometries [15]. Instanton theory over these later spaces possessing several phenomena related with non-compactness (e.g. existence of four dimensional moduli spaces isometric to the original space) is well-known due to the important paper of Kronheimer and Nakajima [27] in which a full ADHM construction was established. The existence of this construction is in some sense not surprising because the original ADHM construction was designed for the flat $\mathbb{R}^4$ and all ALE spaces arise by an algebro-geometric deformation of the *flat* quotients $\mathbb{C}^2/\Gamma$ where $\Gamma \subset \mathrm{SU}(2)$ are various discrete subgroups [26].

The natural question arises: what about instanton theory over ALF spaces? Unlike the ALE geometries, these are essentially non-flat spaces in the sense above therefore one may expect that instanton theory somewhat deviates from that over the flat $\mathbb{R}^4$. Some general questions have been answered recently [13] and these investigations pointed out that in spite of their more transcendental nature, open Riemannian spaces with ALF asymptotics rather resemble compact four manifolds at least from the point of view of instanton theory. To test this interesting observation more carefully in this paper we work out the simplest moduli space over the multi-Taub–NUT geometry. We will find that this space is five rather than four dimensional and looks like a “collar” of the original manifold supporting the analogy with the compact case. We note that this apparent compactness is related to the existence of a smooth compactification of ALF spaces motivated by L^2 cohomology theory [19].

The paper is organized as follows. In Sect. 2 we quickly summarize two important tools for solving the self-duality equations over an anti-self-dual space: the conformal rescaling method of Jackiw–Nohl–Rebbi [25, 4] as well as the Atiyah–Ward correspondence [5]. Via conformal rescaling one constructs instantons out of positive harmonic functions with at most pointlike singularities and appropriate decay toward infinity while the Atiyah–Ward correspondence establishes a link between instantons and holomorphic vector bundles. Since these methods meet in Penrose’ twistor theory we shall also briefly outline it here.

Then we introduce ALF spaces. Referring to recent results on instanton theory over these spaces [13] we precisely define the class of anti-instantons which are expected to form nice moduli spaces. These solutions must obey two conditions: the so-called *weak holonomy condition* with respect to some smooth flat connection and the *rapid decay condition* (cf. Definition 2.1 here). The former condition, albeit looks like an analytical one, in fact deals with the topology of infinity of the ALF space only [13, Theorem 2.3] meanwhile the latter one controls the fall-off properties

of an anti-instanton hence is indeed analytical in its nature.

In Sect. 3, taking probably the most relevant ALF example namely the multi-Taub–NUT series, following Andreson–Kronheimer–LeBrun [1] and LeBrun [28] we review this geometry with special attention to the description of its generic complex structures as algebraic surfaces in $\mathbb{C}^3$. We also identify a finite number of distinguished complex structures that can be realized as blowups of these algebraic surfaces. The group S^1 acts on these spaces via isometries and this action has isolated fixed points, called NUTs.

Then in Sect. 4 we come to our main results. We shall prove via conformal rescaling that over the multi-Taub–NUT space (M_V, g_V) unframed L^2 moduli spaces of $SU(2)$ anti-instantons obeying both the aforementioned weak holonomy condition with respect to the trivial flat connection ∇_Θ as well as the rapid decay condition are not empty. Since the input in this construction is a positive harmonic function, the key concept here is Grigoryan’s *non-parabolic manifolds* [17, 21] which admit an abundance of *positive minimal Green functions* $G(\cdot, y)$ with arbitrarily prescribed pointlike singularity $y \in M_V$ (cf. Definition 4.1 here). Non-parabolicity deals with the volume growth toward infinity and is closely related to the ALF property of our space as it was observed by Minerbe [29, Theorem 0.1]. These functions provide us with a five real parameter family

$$f_{y,\lambda} = 1 + \lambda G(\cdot, y)$$

of positive harmonic functions hence that of non-gauge equivalent irreducible $SU(2)$ -anti-self-dual connections of unital energy on the positive chiral spinor bundle—parameterized by a point $y \in M_V$ and a “concentration parameter” $\lambda \in (0, +\infty)$. These solutions obey the weak holonomy condition with respect to the trivial flat connection as well as decay rapidly. This way we obtain a familiar “collar” for the original manifold (cf. Theorem 4.1 here). Since we know from [13, Theorem 3.2] that the unital energy unframed moduli space of these irreducible solutions is a smooth five dimensional manifold it follows that this collar in fact realizes an open subset of it (or one of its connected component) and describes the regime of concentrated unital energy anti-instantons with trivial holonomy at infinity.

Then we proceed further and construct additional anti-instantons in the “centerless” direction i.e., when the concentration parameter λ becomes infinite. These solutions just correspond to pure Green functions. However in this limit anti-instantons corresponding to the NUTs are reducible to $U(1)$ consequently it is natural to talk about (a connected component of) the unframed moduli space $\widehat{\mathcal{M}}(1, \Theta)$ including these reducible solutions as well. Therefore we will find in particular that this extended moduli space is not a manifold anymore rather admits usual singularities in the reducible points moreover that our moduli space itself admits a singular fibration

$$\Phi : \widehat{\mathcal{M}}(1, \Theta) \longrightarrow \mathbb{R}^3$$

whose generic fibers are 2 dimensional open while the singular ones are 1 dimensional semi-open balls (cf. Theorem 4.2 here). The aforementioned anti-instantons with infinite concentration parameter $\lambda = +\infty$ are represented in the generic case by the centers of these open 2-balls meanwhile in the singular case by the closed end of these semi-open 1-balls. For a given generic fiber the $\lambda = \text{const.} < +\infty$ circles around the center describe anti-instantons “centered” about points in (M_V, g_V) along the same isometry orbit and with fixed finite concentration parameter. The singular fibers correspond to the fixed points of the isometric S^1 -action hence in a given singular fiber there is a unique anti-instanton with $\lambda = \text{const.} \leq +\infty$. In other words, all “centerless” anti-instantons stemming from pure Green functions with pointlike singularities along a given isometry orbit are gauge equivalent.

We will also be able to demonstrate that similar moduli spaces with arbitrary high energy and trivial holonomy at infinity are non-empty according to previous expectations.

For a comparison we also construct the moduli space of rapidly decaying unital energy $SU(2)$ instantons with trivial holonomy in infinity over the flat $\mathbb{R}^3 \times S^1$, too (cf. Theorem 4.4 here).

In the last section, Sect. 5 we round things off by proving, as already mentioned, that our construction is complete in the sense that at least a whole connected component of the basic moduli space emerges this way. This will require however a quite tedious work.

To this end first we provide a detailed construction of the twistor space of the multi-Taub–NUT geometry together with its real structure *à la* Hitchin [22, 24]. Then we transform our anti-instantons constructed out of harmonic functions into holomorphic vector bundles on the twistor space in the spirit of the Atiyah–Ward correspondence [5]. More precisely, guided by a nice result of Atiyah [2] we also work out a smooth compactification of the twistor space which makes it possible to identify the harmonic functions used to construct anti-instantons via conformal rescaling with elements of certain Ext groups of the compactified twistor space (cf. Theorem 5.1 here). These groups can be used to obtain holomorphic bundles on the twistor space. In particular gluing data of the bundles corresponding to minimal Green functions can be worked out explicitly and one can conclude that the isometric S^1 action induces an isomorphism of these bundles therefore the associated “centerless” anti-instantons indeed must be gauge equivalent. This provides us with the picture of the moduli space mentioned above.

Working with the twistor space also can be viewed as a step toward establishing a general ADHM-like instanton factory [8] which can be used to replace the conformal rescaling method; this is certainly necessary since our technique is not capable to grasp all higher energy moduli spaces.

We conjecture that in fact we have obtained the full moduli space of unital energy, rapidly decaying anti-instantons with trivial holonomy at infinity in this paper i.e., the basic moduli space is connected.

Finally for a comparison we make a comment here on other gravitational instantons of ALG and ALH type introduced by Cherkis and Kapustin. By their definition, these geometries are parabolic *à la* Grigoryan hence one cannot expect the existence of positive minimal Green functions over them and accordingly, a canonical five parameter family of anti-instantons. Indeed, the simplest ALG space is the flat $\mathbb{R}^2 \times T^2$ and it is known that the framed unital energy anti-instanton moduli space here is one dimensional only [7]. Hence in the ALE–ALH hierarchy apparently ALF spaces represent the closest analogues of compact four-manifolds from an instanton theoretic viewpoint. Analogously, in this hierarchy ALF spaces are the only ones whose L^2 cohomological compactification [19] is smooth.

2 Constructing instantons

In this section we quickly summarize two methods to obtain instantons over an anti-self-dual Riemannian manifold: the conformal rescaling method and the Atiyah–Ward correspondence. This two approaches meet each other in twistor theory therefore we also review the basic facts about it here.

Let us begin with the conformal rescaling method which was first used by Jackiw, Nohl and Rebbi to obtain instantons over the flat $\mathbb{R}^4$ [25]. Remember that an $SU(2)$ (anti-)self-dual connection ∇_A over a four dimensional oriented Riemannian manifold (M, g) is a smooth $SU(2)$ connection on any $SU(2)$ rank two complex vector bundle E over M whose curvature satisfies the

(anti-)self-duality equations $*_g F_A = \pm F_A$. The gauge equivalence class $[\nabla_A]$ of this connection is called an *(anti-)instanton* if additionally its energy is finite: $\|F_A\|_{L^2(M,g)} < +\infty$. Note that this assumption on the energy is non-trivial only if (M, g) is non-compact but complete.

Let (M, g) be an oriented Riemannian spin four-manifold, possibly non-compact but complete. Take the Levi–Civita connection ∇ of the metric and lift it to get the corresponding spin connection ∇^s . Referring to the splitting $\mathfrak{so}(4) = \mathfrak{su}(2)^+ \oplus \mathfrak{su}(2)^-$ the projected connection $\nabla_A := p^+(\nabla^s)$, with $p^+ : \mathfrak{so}(4) \rightarrow \mathfrak{su}(2)^+$ being the canonical projection, is an $SU(2)^+$ connection on the $SU(2)^+$ rank two complex vector bundle Σ^+ (the positive chiral spinor bundle).

Assume (M, g) is moreover anti-self-dual i.e., $W_g^+ = 0$. Picking up an everywhere smooth and positive function f on M , one can consider the conformally rescaled metric $\tilde{g} := f^2 g$. The new space $(M, \tilde{g})$ still satisfies $W_{\tilde{g}}^+ = 0$ by conformal invariance of the Weyl-tensor. Its scalar curvature however transforms as $f^3 s_{\tilde{g}} = 6\Delta_g f + f s_g$ where Δ_g refers to the scalar Laplacian on (M, g) . If this new scalar curvature $s_{\tilde{g}}$ happens to vanish over M i.e., the rescaling function satisfies the equation

$$\Delta_g f + \frac{s_g}{6} f = 0 \quad (1)$$

then the corresponding projected connection $\nabla_{\tilde{A}}$, constructed like ∇_A before, satisfies the anti-self-duality equations $*_{\tilde{g}} F_{\tilde{A}} = -F_{\tilde{A}}$ over $(M, \tilde{g})$ by a variant of the Atiyah–Hitchin–Singer theorem [4, Proposition 2.2]. However taking into account the conformal invariance of self-duality, we obtain that $\nabla_{\tilde{A}}$ in fact satisfies the *original* $SU(2)^+$ anti-self-duality equations $*_g F_{\tilde{A}} = -F_{\tilde{A}}$ over (M, g) . Hence we have constructed a solution of the anti-self-duality equation via the *conformal rescaling method*.

This anti-self-dual connection locally can be constructed as follows. Let $U \subset M$ be a coordinate ball and let $(\xi^0, \dots, \xi^3)$ denote the orthonormal frame field which diagonalizes $g|_U$. In this gauge write $\nabla|_U = d + \omega$ for the Levi–Civita connection. The rescaled metric $f^2 g$ has a corresponding local frame field $(f\xi^0, \dots, f\xi^3)$ and Levi–Civita connection $\tilde{\nabla}|_U = d + \tilde{\omega}$. The connection 1-form $\tilde{\omega}$ can be calculated by the aid of the Cartan equation $d(f\xi^i) + \tilde{\omega}_j^i \wedge (f\xi^j) = 0$. An easy computation shows that

$$\tilde{\omega}_j^i = \omega_j^i + ((d \log f)\xi_i)\xi^j - ((d \log f)\xi_j)\xi^i$$

where $(\xi_0, \dots, \xi_3)$ denotes the dual vector field on TU (with respect to the original metric $g|_U$) and $(d \log f)\xi_i = g|_U(\text{grad} f, \xi_i)$. Via $\Lambda^1 M \otimes \mathbb{C} \cong \Sigma^+ \otimes \Sigma^-$ we also obtain a gauge on $\Sigma^+|_U$. Lifting $\tilde{\nabla}$ to the spin bundle then projecting it to the positive side (using the 't Hooft matrices, cf. [12]) and exploiting the identification $\mathfrak{su}(2)^+ \cong \text{Im } \mathbb{H}$ given by $(\sigma_1, \sigma_2, \sigma_3) \mapsto (\mathbf{i}, -\mathbf{j}, -\mathbf{k})$ we obtain in this induced gauge

$$\tilde{A} = A + \frac{1}{2} \text{Im}((d \log f)\boldsymbol{\xi}) \quad (2)$$

with $A := p^+(\omega^s)$ being the projection of the original connection and the quaternion-valued 0-form $d \log f$ and 1-form $\boldsymbol{\xi}$ in the second term are defined respectively as follows:

$$d \log f := d \log f(-\xi_0 + \xi_1 \mathbf{i} + \xi_2 \mathbf{j} + \xi_3 \mathbf{k}), \quad \boldsymbol{\xi} := \xi^0 + \xi^1 \mathbf{i} + \xi^2 \mathbf{j} + \xi^3 \mathbf{k}.$$

Writing $a|_U := \frac{1}{2} \text{Im}((d \log f)\boldsymbol{\xi})$ for the only term involving f , it is clear that it extends over M as an $\text{Im } \mathbb{H}$ -valued 1-form i.e., one finds

$$a \in C^\infty(\Lambda^1 M \otimes \text{End } \Sigma^+) \quad (3)$$

and it is insensitive for an overall rescaling of f by a non-zero real constant. This implies that from the point of view of our anti-instanton factory, we can regard two conformal scaling functions identical if they differ by an overall rescaling only. Moreover note that the new connection takes the form $\nabla_{\tilde{A}} = \nabla_{A+a}$ i.e., this whole conformal rescaling procedure can be regarded as an anti-self-dual perturbation of the original connection. For an explicit application of this method we refer to [25] over the flat $\mathbb{R}^4$ and to [12] in case of the multi-Taub–NUT space.

If (M, g) is compact then this machinery provides us with an anti-instanton on Σ^+ . If the space is non-compact but complete and ∇_A has finite energy (which can be achieved if the metric decays sufficiently fast to the flat one at infinity as we have seen) then if the perturbation a also decays fast i.e., f is sufficiently bounded at infinity then $\nabla_{\tilde{A}}$ has still finite energy consequently we obtain an anti-instanton on the same bundle Σ^+ over (M, g) .

An important particular case is when the original space (M, g) has already vanishing scalar curvature. In this case (1) cuts down to the ordinary Laplace equation $\Delta_g f = 0$ and the original connection ∇_A is already anti-self-dual corresponding to the trivial solution $f = 1$.

However it is clear by the maximum principle that in many situations there are no non-trivial everywhere smooth bounded solutions to (1) therefore this method is apparently vacuous. Fortunately we can modify f to possess mild singularities over M . At this level of generality by a “mild singularity” we mean the following:

- (i) There are finitely many points $\{y_1, \dots, y_k\} \subset M$ such that f diverges in y_i for all $i = 1, 2, \dots, k$;
- (ii) The local energy of the rescaled connection around each point is finite i.e., for U_i a neighbourhood of y_i one has

$$\|F_{\tilde{A}}\|_{L^2(U_i \setminus \{y_i\}, g)} < +\infty$$

for all $i = 1, 2, \dots, k$. (In terms of f this condition simply controls its behaviour about y_i .)

Under these circumstances Uhlenbeck’s theorem [32] allows us to remove the singularities from $\nabla_{\tilde{A}}$ introduced by f . Therefore if f is a positive solution to (1) with such singularities then the resulting anti-self-dual connection will be smooth again and corresponds to a non-trivial anti-instanton over a bundle E if M is compact. This $SU(2)^+$ bundle may be no more isomorphic to Σ^+ as a consequence of the allowed singularities of the scaling function. In fact its topological type is determined by the singularity structure of f . In case of a non-compact complete base geometry if in addition f is sufficiently bounded at infinity then the resulting anti-self-dual connection has finite energy i.e., describes an anti-instanton on the same bundle Σ^+ , regardless how many singularities f possesses. This is because in fact over a non-compact four-manifold all $SU(2)^+$ bundles, including Σ^+ , are isomorphic to the trivial one. The significance of the singularity structure of scaling functions in instanton theory therefore becomes transparent.

We move on and recall twistor theory of anti-self-dual geometries and in particular the Atiyah–Ward construction [5] which is another powerful way to solve the (anti-)self-duality equations at least in principle.

Let us recall the general theory [6, Chapter 13]. Let (M, g) be an oriented Riemannian four-manifold and consider the projectivization $Z := P(\Sigma^+)$ of the positive chiral spinor bundle Σ^+ on M (this projectivization is well-defined even if M is not spin). Clearly, Z admits a fiber bundle structure $p : Z \rightarrow M$ with $\mathbb{CP}^1$ ’s as fibers. A fiber F_x over $x \in M$ can be viewed as the parameter space of orthogonal complex structures on $T_x M$ whose induced orientations agree with that of M . Using the Levi–Civita connection on (M, g) we can split the tangent space

$T_z Z$ at any point $z \in Z$; writing $z = (x, p)$ with $x \in M$ and $p \in F_x$ one endows the horizontal subspace $H_{(x,p)} \cong T_x M$ of $T_{(x,p)} Z$ with the complex structure J_p given by $p \in F_x$ itself meanwhile the vertical subspace $V_{(x,p)} \cong T_p F_x$ of $T_{(x,p)} Z$ is equipped with the standard complex structure I coming from $F_x \cong \mathbb{C}P^1$. Therefore $T_{(x,p)} Z$ carries a complex structure (J_p, I) and Z as a real six-manifold gets possess an almost complex structure. A basic theorem of Penrose [31] or Atiyah, Hitchin and Singer [4, Theorem 4.1][6, Theorem 13.46] states that this complex structure is integrable i.e., Z is a complex three-manifold if and only if (M, g) is anti-self-dual. The space Z is called the *twistor space* of (M, g) . The complex structure in fact depends on the conformal class of the metric only.

The basic holomorphic structure of Z looks like as follows [6, Chapter 13]:

- (i) For all $y \in M$ the fibers $F_y \subset Z$ represent *holomorphic lines* $Y \subset Z$ with normal bundles isomorphic to $H \oplus H$ where H is the usual line bundle H with $\langle c_1(H), [Y] \rangle = 1$ on $Y \cong \mathbb{C}P^1$;
- (ii) The antipodal map on Y induces a *real structure* on Z i.e., an anti-holomorphic involution $\tau : Z \rightarrow Z$ satisfying $\tau^2 = \text{Id}_Z$;
- (iii) The space of all holomorphic lines in Z near the fibers Y form a locally complete family which is a complex manifold $M^\mathbb{C}$ of complex dimension four and the fibers Y among them are distinguished by the property that the real structure fixes them, hence are called *real lines*.

A remarkable property of twistor spaces is that they allow one to solve certain linear or non-linear field equations over (M, g) . These equations are of great importance in physics. We shall also quickly review these constructions.

1. The linear field equation relevant to us here is the *conformal scalar Laplace equation* (1). We shall denote by H_Z the holomorphic line bundle on the whole Z such that for any fiber the restriction $H_Z|_Y$ is the standard line bundle H on $Y \cong \mathbb{C}P^1$. We define the bundles H_Z^k on Z with $k \in \mathbb{Z}$ analogously. For notational convenience write $\mathcal{O}_Z(k)$ for the sheaf $\mathcal{O}(H_Z^k)$. Recall (cf. e.g. [23]) that complex-valued real analytic functions satisfying (1) on open subsets $(U, g|_U) \subset (M, g)$ are in one-to-one correspondence with elements of the sheaf cohomology group $H^1(p^{-1}(U); \mathcal{O}_{p^{-1}(U)}(-2))$ i.e., there is a natural isomorphism

$$T : H^1(p^{-1}(U); \mathcal{O}_{p^{-1}(U)}(-2)) \cong \text{Ker } \Delta_g|_U$$

called the *Penrose transform*: given an element $\varphi \in H^1(p^{-1}(U); \mathcal{O}_{p^{-1}(U)}(-2))$ and a real line $Y \subset p^{-1}(U)$ one can take the restriction $\varphi|_Y \in H^1(Y; \mathcal{O}_Y(-2)) \cong \mathbb{C}$. This gives rise to a complex-valued function $f(y) := \varphi|_Y$ consequently one has a map

$$T : H^1(p^{-1}(U); \mathcal{O}_{p^{-1}(U)}(-2)) \longrightarrow C^\infty(U, \mathbb{C}).$$

It turns out that f is a solution to (1) on $U \subset M$ and all local solutions arise this way.

2. The non-linear *(anti-)self-duality equations* of Yang–Mills theory, also can be treated in twistor theory (cf. e.g. [4]): solutions of the $\text{SU}(2)$ anti-self-duality equations over (M, g) with appropriate orientation can be converted into certain holomorphic vector bundles on Z as follows. There is a one-to-one correspondence of gauge equivalence classes of anti-self-dual $\text{SU}(2)$ connections on a rank two complex vector bundle E on M and isomorphism classes of rank two complex vector bundles F on Z such that

- (i) F is holomorphic;

- (ii) For any real line $Y \subset Z$ the restricted bundle $F|_Y$ is holomorphically trivial;
- (iii) There exists a conjugate linear map $\tilde{\tau} : F \rightarrow F$ lying over the real structure $\tau : Z \rightarrow Z$ and satisfies $\tilde{\tau}^2 = -\text{Id}_F$.

If ∇_A is irreducible and Z (that is, M) is compact then F is a stable holomorphic bundle on Z and in this case $\tilde{\tau}$ is unique up to scalar multiplication.

This construction, called the *Atiyah–Ward correspondence*, goes as follows. Take an $\text{SU}(2)$ vector bundle E and an anti-self-dual connection on it over (M, g) . Let $F := p^*E$ be a C^∞ bundle; by anti-self-duality the curvature of the lifted connection $p^*\nabla_A$ will be of $(1, 1)$ -type hence F will be holomorphic. One checks that properties (ii) and (iii) are also satisfied. Conversely, take a holomorphic bundle F satisfying (ii) and (iii). Define E_y to be the space of holomorphic sections over $F|_Y$ (this space is two complex dimensional via (ii)). Property (iii) induces a symplectic structure on F consequently we can take the corresponding unique unitary connection ∇ on F . By uniqueness and (ii) this connection is the trivial connection on each $F|_Y$ hence is of the form $p^*\nabla_A$ with a connection ∇_A on E . Since the curvature of this connection is of $(1, 1)$ -type with respect to all complex structures on $T_y M$ for all $y \in M$ it follows that it is anti-self-dual.

3. Another non-linear equation, namely the *(Riemannian) self-dual vacuum Einstein equations* of general relativity can also be naturally adjusted to twistor theory, known as *Penrose’ non-linear graviton construction* [31, 22]. Assume here that (M, g) is moreover simply connected and Ricci-flat. Then it follows that the induced connection on Σ^+ is the trivial flat one and Z can be retracted onto its particular fiber F_y by parallel transport. Therefore topologically $Z \cong M \times S^2$ in this case. The associated twistor space of (M, g) has in addition to the basic holomorphic properties above the following:

- (i) There is a holomorphic fibration $\pi : Z \rightarrow Y$ over a particular line $Y \cong \mathbb{CP}^1$ with fibers being the original space M . Consequently $H_Z^k \cong \pi^* H^k$ for all $k \in \mathbb{Z}$;
- (ii) Since the complex normal bundle N_Y of each Y is isomorphic to $H \oplus H$ it follows that the canonical bundle K_Z of Z is isomorphic to $\pi^* H^{-4}$.

Conversely, it can be shown [6, Chapter 13] that a complex 3-space Z having these properties encodes the conformal class of a Ricci-flat, anti-self-dual simply connected four-space (M, g) . Indeed, the complexified space $M^\mathbb{C}$ carries a natural conformal structure by declaring two points $Y', Y'' \in M^\mathbb{C}$ to be null-separated if and only if for the corresponding lines $Y' \cap Y'' \neq \emptyset$ in Z . Restricting this conformal structure to the real lines parameterized by $M \subset M^\mathbb{C}$ one comes up with the conformal class of an anti-self-dual Ricci-flat structure (M, g) . The particular metric emerges via an appropriate isomorphism $K_Z \cong \pi^* H^{-4}$.

We have encountered two techniques for constructing (anti)-instantons over an anti-self-dual space. The link between them is an observation of Atiyah [2] which relates solutions to (1) to holomorphic vector bundles on the twistor space. We will work out this in detail in Sect. 5.

In the rest of this section we introduce a special class of four-manifolds: suppose from now on that (M, g) is a four dimensional *asymptotically locally flat* (ALF) space as it was defined in [13]. By a recent powerful theorem of Minerbe [29, Theorem 0.1] these four dimensional geometries at least in the connected, geodesically complete, hyper-Kähler case can also be characterized by requiring their curvature to be L^2 with respect to the measure $\frac{r^4}{\text{Vol}_g(B_r^4(x))} dV_g$ as well as prescribing their asymptotical volume growth to be $O(r^\nu)$ with $3 \leq \nu < 4$.

Framed moduli spaces of certain “admissible” $SU(2)$ (anti-)instantons over these geometries have been investigated recently [13]. By “admissible” we mean the following. Topologically, an ALF space (M, g) (with a single end) admits a decomposition $M = K \cup W$ where K is a compact interior space and W is an end or neck homeomorphic to $N \times \mathbb{R}^+$ where $\pi_N : N \rightarrow B_\infty$ is a connected, compact, oriented three-manifold fibered over a compact Riemann surface B_∞ with circle fibers $F \cong S^1$. Note that for any $x \in M$ and sufficiently large $R > 0$ W is homeomorphic to $M \setminus B_R^4(x)$ where $B_R^4(x)$ is a geodesic ball about $x \in M$ with radius $R > 0$.

Definition 2.1. *Let (M, g) be an ALF four-manifold. Take an arbitrary smooth, finite energy $SU(2)$ connection ∇_A on a (necessarily trivial) $SU(2)$ vector bundle E over M . This connection is said to be admissible if it satisfies two conditions (cf. [13]):*

- (i) *The first is called the weak holonomy condition and says that to ∇_A there exists a smooth flat $SU(2)$ connection $\nabla_\Gamma|_W$ on $E|_W$ along the end $W \subset M$ and a constant $c_1 = c_1(g) > 0$, independent of $R > 0$, such that in any smooth gauge on $M \setminus B_R^4(x)$ the inequality*

$$\|A - \Gamma\|_{L^2_{1,\Gamma}(M \setminus B_R^4(x), g)} \leq c_1 \|F_A\|_{L^2(M \setminus B_R^4(x), g)}$$

holds along the neck $M \setminus B_R^4(x)$;

- (ii) *The second condition requires ∇_A to decay rapidly at infinity i.e.,*

$$\lim_{R \rightarrow \infty} \sqrt{R} \|F_A\|_{L^2(M \setminus B_R^4(x), g)} = 0.$$

The first condition ensures us that the connection has a well-defined holonomy at infinity given by $\nabla_\Gamma|_W$; the second condition regulates how fast the finite energy connection decays to this flat connection and also can be reformulated (cf. [13]) by saying that F_A belongs to a weighted Sobolev space with weight $\delta = \frac{1}{2}$ i.e., $F_A \in L^2_{\frac{1}{2}}(\Lambda^2 M \otimes \text{End } E)$. Note that this second condition is stronger than the finite energy requirement on ∇_A . It was demonstrated in [13] that irreducible $SU(2)$ anti-instantons satisfying conditions (i) and (ii) with a fixed limiting flat connection $\nabla_\Gamma|_W$ and energy e form framed moduli spaces $\mathcal{M}(e, \Gamma)$ which are smooth finite dimensional manifolds. Their dimensions have also been calculated by the aid of a dimension formula.

Now take an ALF space (M, g) with a fixed orientation whose curvature satisfies $W_g^+ = 0$ and $s_g = 0$ therefore the Atiyah–Hitchin–Singer theorem applies and the projected Levi–Civita connection ∇_A is an anti-self-dual connection on Σ^+ . By an appropriate curvature decay imposed on the metric (which is incorporated into the precise definition of an ALF space, cf. [13]) this connection has not only finite energy but also decays rapidly as in (ii) of Definition 2.1; moreover if additionally it satisfies the weak holonomy condition (i) in Definition 2.1 then it gives rise to an anti-instanton on Σ^+ in the sense above. Performing a conformal rescaling of the metric we obtain another anti-self-dual connection $\nabla_{\tilde{A}}$ on Σ^+ . Since it is of the form ∇_{A+a} with a as in (3) except finitely many points, its curvature reads as $F_{\tilde{A}} = F_{A+a} = F_A + d_A a + a \wedge a$ consequently for its energy

$$\|F_{\tilde{A}}\|_{L^2_{\frac{1}{2}}(M, g)} \leq \|F_A\|_{L^2_{\frac{1}{2}}(M, g)} + \|d_A a\|_{L^2_{\frac{1}{2}}(M, g)} + c_2 \|a\|_{L^2_{\frac{1}{2}, 1, \Gamma}(M, g)}^2 \quad (4)$$

holds with a constant $c_2 = c_2(g) > 0$. We conclude that the regularity and decay properties of the perturbed connection is determined by the perturbation term. More precisely, the condition

$$a \in L^2_{\frac{1}{2}, 1, \Gamma}(\Lambda^1 M \otimes \text{End } \Sigma^+), \quad \tilde{A} \in L^2_{loc, 1, \tilde{A}}(\Lambda^1 M \otimes \text{End } \Sigma^+), \quad F_{\tilde{A}} \in L^2_{loc}(\Lambda^- M \otimes \text{End } \Sigma^+) \quad (5)$$

ensures us that a affects neither the asymptotics of ∇_A or its fall-off and any pointlike singularities are removable. Therefore if ∇_A is smooth and obeys both the weak holonomy and the rapid decay conditions above then $\nabla_{\tilde{A}} = \nabla_{A+a}$ will be also smooth, decay rapidly and converges to the the same limiting flat connection. More precisely one can show that the energy $\tilde{e}$ of $\nabla_{\tilde{A}}$ can differ from that of ∇_A by a non-negative integer only [13]. That is, $[\nabla_A] \in \mathcal{M}(e, \Gamma)$ implies $[\nabla_{\tilde{A}}] \in \mathcal{M}(\tilde{e}, \Gamma)$ and $\tilde{e} - e \in \{0, 1, 2, \dots\}$. In fact as one expects, the difference $\tilde{e} - e$ is governed by the singularity structure of the perturbation i.e., of the scaling function f in (1).

In the next sections we will apply the conformal rescaling technique and twistor theory to understand the moduli space over the multi-Taub–NUT geometries of unital energy $SU(2)$ anti-instantons decaying rapidly and obeying the weak holonomy condition with respect to the trivial connection ∇_Θ .

3 Review of the multi-Taub–NUT geometry

In this section we take a closer look of the multi-Taub–NUT spaces following [1, 28]. Consider $s > 0$ distinct points $q_1, \dots, q_s \in \mathbb{R}^3$. We will construct a four-manifold M_V and a smooth map $\pi : M_V \rightarrow \mathbb{R}^3$ such that $\pi^{-1}(q_j)$ is a point for all $j = 1, 2, \dots, s$ but $\pi^{-1}(x) \cong S^1$ for all other points $x \in \mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$. To begin with, let $\pi : U_V \rightarrow \mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$ be the principal S^1 bundle whose Chern class is -1 when restricted to a small sphere of radius $r_j < \min_{k \neq j} |q_j - q_k|$ around q_j . Thus $\pi^{-1}(B_{r_j}^3(q_j))$ is diffeomorphic to the punctured ball $B_j^4 \setminus \{0\} \subset \mathbb{R}^4$ in a manner such that the S^1 action becomes the action of $S^1 \subset \mathbb{C}$ on $\mathbb{C}^2 \cong \mathbb{R}^4$ by scalar multiplication. We then define

$$M_V := (U_V \sqcup B_1^4 \sqcup \dots \sqcup B_s^4) / \sim$$

where $\sim$ means that $B_j^4 \setminus \{0\}$ is identified with $\pi^{-1}(B_{r_j}^3(q_j))$. The map $\pi : U_V \rightarrow \mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$ extends to a smooth map $\pi : M_V \rightarrow \mathbb{R}^3$. Note that there is an S^1 action on M_V whose fixed points with index -1 are exactly the q_j 's—called *NUTs*. If ℓ_{ij} is a straight line segment joining q_i with q_j then $\pi^{-1}(\ell_{ij}) \subset M_V$ is a smoothly embedded 2-sphere whose self-intersection number is -2 . The 2-spheres $\pi^{-1}(\ell_{j,j+1})$ with $j = 1, 2, \dots, s-1$ are attached together according to the A_{s-1} Dynkin diagram and generate $H^2(M_V, \mathbb{Z}) \cong \mathbb{Z}^{s-1}$.

Take a real number $c > 0$ and let $V : \mathbb{R}^3 \setminus \{q_1, \dots, q_s\} \rightarrow \mathbb{R}^+$ be defined by

$$V(x) := c + \frac{1}{2} \sum_{j=1}^s \frac{1}{|x - q_j|}.$$

In this paper we will suppose $c = 1$. Then $\Delta V = 0$ that is, it is a positive harmonic function on $\mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$. The cohomology class $[\frac{1}{2\pi}\pi^*(\ast_3 dV)] \in H^2(U_V, \mathbb{R})$ is the first Chern class of the bundle $\pi : U_V \rightarrow \mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$. Consequently there is a connection ∇ on U_V whose curvature is $\ast_3 dV$. More precisely, let ω be the imaginary-valued connection 1-form on U_V so that $\mathbf{i}\pi^*(\ast_3 dV) = F_\omega = d\omega$. The form ω is unique up to gauge transformation since $\mathbb{R}^3 \setminus \{q_1, \dots, q_s\}$ is simply connected. If we introduce a coordiante system (x^1, x^2, x^3, τ) where x^i are Cartesian coordinates on $\mathbb{R}^3$ and $\tau \in [0, 2\pi)$ parameterizes the circles on U_V then we write $\frac{1}{\mathbf{i}}\omega = d\tau + \pi^*\alpha$ with $\alpha \in C^\infty(\Lambda^1(\mathbb{R}^3 \setminus \{q_1, \dots, q_s\}))$ a real-valued 1-form. The *multi-Taub–NUT metric* $g_V|_{U_V}$ on U_V is defined to be

$$ds^2 = V\pi^*((dx^1)^2 + (dx^2)^2 + (dx^3)^2) + \frac{1}{V}(d\tau + \pi^*\alpha)^2. \quad (6)$$

The space $(U_V, g_V|_{U_V})$ extends smoothly across the fixed points of the S^1 action: write $V(x) = V_j(x) + f(x)$ with $V_j(x) := \frac{1}{2|x-q_j|}$ and f a smooth function around q_j . Let ω_j be the connection whose curvature is $\pi^*(\ast_3 dV_j)$ then we obtain a decomposition

$$ds^2 = V_j \pi^*((dx^1)^2 + (dx^2)^2 + (dx^3)^2) + \frac{1}{V_j} (d\tau + \pi^* \alpha_j)^2 + h$$

where h is a smooth symmetric bilinear form around p_j induced by f . Moreover the first term is just the 1-Eguchi–Hanson metric on $\mathbb{R}^4$ which is known to be isometric to the standard flat metric on $\mathbb{R}^4$ hence extends over p_j .

We come therefore up with a Riemannian manifold (M_V, g_V) with an isometric S^1 action whose fixed points with index -1 are $\pi^{-1}(q_j)$. It is moreover complete with a single ALF end, is anti-self-dual since $d\alpha = \ast_3 dV$ and in particular is Ricci-flat. Since M_V is simply connected, it follows that it is hyper-Kähler as well i.e., there is an entire 2-sphere's worth of complex structures for which g_V is a Kähler metric.

To display these complex structures explicitly at least on U_V , let e_1, e_2, e_3 be an orthonormal basis in $\mathbb{R}^3$. Consider these as constant vector fields on $\mathbb{R}^3$ and let $\hat{e}_1, \hat{e}_2, \hat{e}_3$ be their horizontal lifts to U_V via the connection, as well as let $\hat{e}_0$ be the generator of the S^1 -action. Then

$$(\xi_0, \xi_1, \xi_2, \xi_3) := \left(\sqrt{V} \hat{e}_0, \frac{1}{\sqrt{V}} \hat{e}_1, \frac{1}{\sqrt{V}} \hat{e}_2, \frac{1}{\sqrt{V}} \hat{e}_3 \right) \quad (7)$$

is an orthonormal frame on (U_V, g) which we define to be oriented. Relative to this frame we take the almost complex structure

$$J_{e_1} := \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

It is parallel hence is integrable and it depends on the choice of e_1 . Of course it extends over the whole (M_V, g_V) which we continue to denote by J_{e_1} . By the aid of this picture we see that all possible orthogonal complex structures with compatible orientation on (M_V, g_V) are parameterized by a $\mathbb{C}P^1$ and the situation can be described as a holomorphic fibration: the fiber over $e_1 \in \mathbb{C}P^1$ is the complex manifold (M_V, J_{e_1}) in accord with the general theory outlined in Sect. 2. Moreover if the direction of any of the straight line segments ℓ_{ij} coincides with e_1 then the corresponding 2-sphere $\pi^{-1}(\ell_{ij})$ will represent a holomorphic curve with self-intersection number -2 in (M_V, J_{e_1}) . There are at most $2\binom{s}{2} = s(s-1)$ such exceptional complex structures. For a generic complex structure however, there are no holomorphic projective lines.

These complex structures on the whole (M_V, g_V) can be described in a rather explicit algebraic way as follows (cf. [1, 28]).

Lemma 3.1. *Fix a generic configuration of points $q_1, \dots, q_s$ in $\mathbb{R}^3$ and consider the associated space (M_V, g_V) . Take a direction e_1 in $\mathbb{R}^3$. Assume that e_1 is not parallel with any line segments ℓ_{ij} joining q_i with q_j . Then (M_V, J_{e_1}) is biholomorphic to the algebraic surface $X \subset \mathbb{C}^3$ given by*

$$xy - (z - p_1) \dots (z - p_s) = 0 \quad (8)$$

with $(x, y, z) \in \mathbb{C}^3$ and fixed distinct complex numbers $p_1, \dots, p_s \in \mathbb{C}$.

The S^1 -action on X is given by $(x, y, z, \tau) \mapsto (xe^{i\tau}, ye^{-i\tau}, z)$ with fixed points $(0, 0, p_j) \in X$ of index -1 for all $j = 1, \dots, s$. Under this biholomorphism the fixed point $\pi_V^{-1}(q_j) \in M_V$ of the S^1 -action on M_V correspond to the fixed point $(0, 0, p_j) \in X$ with the same index for all $j = 1, \dots, s$ consequently with this S^1 -action on X we obtain an isomorphism of complex manifolds with an S^1 -action.

If e_1 is parallel with any ℓ_{ij} then (M_V, J_{e_1}) is biholomorphic to the complex surface $\tilde{X}$ which arises by taking $p_i = p_j$ for fixed i and j in (8) and blowing up the resulting singular surface $X^* \subset \mathbb{C}^3$ in $(0, 0, p_j) \in X^*$.

As a real oriented four-manifold $\tilde{X}$ is diffeomorphic to X .

Proof. By genericity we assume that the straight line segments $\ell_{ij} \subset \mathbb{R}^3$ with $i, j = 1, 2, \dots, s$ are mutually non-parallel. First take a direction e_1 in $\mathbb{R}^3$ which is not parallel with any ℓ_{ij} . Let $(e_2 e_3) \subset \mathbb{R}^3$ be the oriented plane passing through the origin, perpendicular to e_1 and identify it with the complex plane with its standard orientation: $(e_2 e_3) \cong \mathbb{C}$ such that $e_2 \mapsto 1$ and $e_3 \mapsto i$. Let $p_{e_1} : \mathbb{R}^3 \rightarrow \mathbb{C}$ be the projection along e_1 which induces $P_{e_1} := p_{e_1} \circ \pi : M_V \rightarrow \mathbb{C}$, the projection from M_V . Then P_{e_1} is a holomorphic map from (M_V, J_{e_1}) to $\mathbb{C}$. For a point $a \in M_V$ define a complex coordinate by $z := P_{e_1}(a) \in \mathbb{C}$. In particular for a NUT $q_j \in \mathbb{R}^3$ put $p_j := P_{e_1}(\pi^{-1}(q_j)) = p_{e_1}(q_j) \in \mathbb{C}$ for all $j = 1, \dots, s$ and also call them NUTs.

Next we analyze the preimage $P_{e_1}^{-1}(z) \subset M_V$. If $z \neq p_j \in \mathbb{C}$ then $P_{e_1}^{-1}(z)$ is holomorphically isomorphic to an infinite cylinder $\mathbb{C}^* := \{(x, y) \in \mathbb{C}^2 \mid xy = 1\}$ in (M_V, J_{e_1}) , since the directions $\hat{e}_0$ and $\hat{e}_1$ span a holomorphic line for the complex structure J_{e_1} . On the other hand $P_{e_1}^{-1}(p_j)$ is homeomorphic to two complex affine lines intersecting in one point. In particular we have a model like $\{(x, y) \in \mathbb{C}^2 \mid xy = 0\}$ for $P_{e_1}^{-1}(p_j)$. Since the complex structure on the fibers is independent of z , the above two cases can be managed together into a global equation by writing $xy = c(z)$ where $c(z)$ is a polynomial in the variable z on $\mathbb{C}$ which vanishes exactly in the points $p_1, \dots, p_s \in \mathbb{C}$. Therefore we put

$$c(z) := (z - p_1) \dots (z - p_s)$$

to obtain the desired biholomorphism between (M_V, J_{e_1}) and X given by (8).

The statement about the S^1 -action is clear.

If e_1 happens to be parallel with any ℓ_{ij} connecting $q_i \in \mathbb{R}^3$ and $q_j \in \mathbb{R}^3$ then we find $p_{e_1}(q_i) = p_i = p_{e_1}(q_j) = p_j \in \mathbb{C}$ hence two roots coincide in (8). However now $P_{e_1}^{-1}(p_i) = P_{e_1}^{-1}(p_j)$ consists of not only two complex affine lines as above but in addition the projective line $\pi^{-1}(\ell_{ij})$ and the two complex affine lines do not intersect each other rather each individual line hits $\pi^{-1}(\ell_{ij})$ in one point respectively. These two points are distinct. The situation can be modeled on $\{(x, y, [u : v]) \in \mathbb{C}^2 \times \mathbb{C}P^1 \mid xy = 0, xv = yu\}$. Therefore to obtain the resulting smooth complex surface we have to take a singular curve $X^* \subset \mathbb{C}^3$ represented by (8) with $p_i = p_j$ and blow it up at $(0, 0, p_i) = (0, 0, p_j) \in X^*$ to obtain $\tilde{X}$.

It readily follows that as a real oriented four-manifold, $\tilde{X}$ is diffeomorphic to X since both are just the original M_V equipped with different complex structures. $\diamond$

Remark. Incidentally we note that the above definition of the coordinate z depends on the identification of the oriented plane perpendicular to e_1 with $\mathbb{C}$, i.e. on the choice of a vector e_2 in the tangent space of $\mathbb{C}P^1$ at e_1 . For all such choices, the above procedure yields a complex number. As $T\mathbb{C}P^1 \cong H^2$, this means that globally over $\mathbb{C}P^1$ the coordinate z must be regarded as a section of the line bundle H^2 . This fact will play an important role in constructing the twistor space, cf. Sect. 5.

4 Moduli spaces with trivial holonomy

We already know that a multi-Taub–NUT space (M_V, g_V) is anti-self-dual and Ricci-flat. Consequently for its curvature $W_{g_V}^+ = 0$ and $s_{g_V} = 0$ holds and the Atiyah–Hitchin–Singer theorem applies and we can construct $SU(2)^+$ (anti-)self-dual connections on the (trivial) positive chiral spinor bundle Σ^+ . If we fix an orientation coming from any complex structure in the hyper-Kähler family, these connections will be anti-self-dual. As we have seen in Section 2, this approach amounts to find positive solutions of the Laplace equation $\Delta_{g_V} f = 0$ with a finite number of mild singularities and appropriate bounds at infinity over the original manifold (M_V, g_V) . This program partially was carried out in [12, 10] and will be stretched to its limits here.

An abundance of such harmonic functions are provided by minimal positive Green functions therefore we make a short digression on them following [21].

Definition 4.1. *Let (M, g) be a non-compact complete Riemannian manifold of dimension $m \geq 2$ and $y \in M$ be a point. A function $G(\cdot, y) : M \setminus \{y\} \rightarrow \mathbb{R}$ is called the minimal positive Green function concentrated at $y \in M$ if it has the following properties:*

- (i) *It satisfies $\Delta_g G(\cdot, y) = \delta_y$ in the sense of distributions on (M, g) ;*
- (ii) *If $r := d(\cdot, y)$ is the distance from y on (M, g) then $G(\cdot, y)$ tends to zero as $r \rightarrow +\infty$ and $G(\cdot, y)$ is $O(r^{2-m})$ and $|\nabla_y G(\cdot, y)|_g$ is $O(r^{1-m})$ for $m \geq 3$ as well as $G(\cdot, y)$ is $O(\log r)$ and $|\nabla_y G(\cdot, y)|_g$ is $O(r^{-1})$ for $m = 2$, if $r \rightarrow 0$;*
- (iii) *Moreover $0 < G(\cdot, y) \leq G'(\cdot, y)$ for any other positive Green function with the same singularity.*

This function is obviously unique and is characterized by these properties.

Such functions may be constructed as follows. Take $y \in M$ and $\Omega \Subset M$ an open subset with smooth boundary and $\bar{\Omega}$ compact such that $y \in \Omega$. Let $G_\Omega(\cdot, y)$ be the unique Green function on Ω satisfying

$$\Delta_g G_\Omega(\cdot, y) = \delta_y \quad \text{and} \quad G_\Omega(\cdot, y)|_{\partial\bar{\Omega}} = 0.$$

By the maximum principle we can assume that $G_\Omega(\cdot, y)$ is positive on $\Omega \setminus \{y\}$ and, if we extend $G_\Omega(\cdot, y)$ by zero outside Ω , then $G_\Omega(\cdot, y) \leq G_{\Omega'}(\cdot, y)$ whenever $\Omega \subseteq \Omega'$. Consequently setting

$$G(\cdot, y) := \sup_{\Omega} G_\Omega(\cdot, y)$$

it follows that if this function exists then it is the minimal positive Green function with singularity in $y \in M$. One can prove [21, Theorem 8.1] that either minimal positive Green functions with singularity $y \in M$ exist over M , uniformly in y , or such functions do not exist at all. In the former case (M, g) is called *non-parabolic*, while in the later case called *parabolic* [17].

Varopoulos showed [33] that if the Ricci curvature of (M, g) is non-negative and for some point $x \in M$ with its geodesic ball of radius r

$$\int_1^{+\infty} \frac{r}{\text{Vol}_g(B_r^4(x))} dr < +\infty$$

then (M, g) is non-parabolic. For the Ricci-flat multi-Taub–NUT space (M_V, g_V) we know that

$$\text{Vol}_{g_V}(B_r^4(x)) \sim \frac{8\pi^2}{3} r^3$$

demonstrating that it is indeed ALF but still non-parabolic; consequently for all $y \in M_V$ an associated minimal positive Green function $G(\cdot, y)$ exists. In fact they were explicitly constructed by Page [30] and the simplest of them also appeared in [12]. Making use of these functions we obtain plenty of positive harmonic functions of the form

$$f_{y,\lambda}(x) := 1 + \lambda G(x, y) \quad (9)$$

with a real constant $\lambda \in (0, +\infty)$.

We claim that these functions have the required properties and can be used to construct finite-energy smooth $SU(2)^+$ anti-instantons over the multi-Taub–NUT space. To check this, we construct the associated five parameter family $\nabla_{\tilde{A}_{y,\lambda}}$ of non-gauge equivalent anti-self-dual $SU(2)^+$ connections on Σ^+ with respect to the orientation on (M_V, g_V) coming from the complex structures and demonstrate first that they are smooth around $y \in M_V$ hence everywhere and secondly that they are admissible hence in particular have finite energy.

Lemma 4.1. *Fix a point $y \in M_V$ and a number $\lambda \in (0, +\infty)$. Take a geodesic ball $B_\varepsilon^4(y) \subset M_V$ around y and write $B_\varepsilon^*(y) := B_\varepsilon^4(y) \setminus \{y\}$ for the punctured ball. Consider the orthonormal frame (γ) on $(B_\varepsilon^4(y), g_V)$. In this gauge plugging (9) into (2) write $\nabla_{\tilde{A}_{y,\lambda}}|_{B_\varepsilon^*(y)} = d + \tilde{A}_{y,\lambda}$ for the resulting anti-self-dual connection.*

There exists an $L^2_{1,\tilde{A}_{y,\lambda}}$ gauge transformation $\gamma : B_\varepsilon^(y) \rightarrow SU(2)^+$ such that the gauge transformed potential $\tilde{A}'_{y,\lambda} := \gamma^{-1}\tilde{A}_{y,\lambda}\gamma + \gamma^{-1}d\gamma$ extends smoothly over $B_\varepsilon^4(y)$ that is the $SU(2)^+$ connection $\nabla_{\tilde{A}_{y,\lambda}}|_{B_\varepsilon^4(y)}$ is smooth on $\Sigma^+|_{B_\varepsilon^4(y)}$.*

Consequently the anti-self-dual connection $\nabla_{\tilde{A}_{y,\lambda}}$ is smooth everywhere on Σ^+ .

Proof. Take an orthonormal frame $(\xi^0, \xi^1, \xi^2, \xi^3)$ over $B_\varepsilon^4(y)$ as in (7) with its associated complex structure J_{e_1} on M_V . Since the multi-Taub–NUT metric is Kähler with respect to this complex structure, there exists a complex coordinate system (z^1, z^2) on $B_\varepsilon^4(y)$ centered in y such that

$$g_V|_{B_\varepsilon^4(y)} = (\delta_{ij} + O(r^2))dz^i d\bar{z}^j$$

i.e., the metric osculates the flat metric in second order. Take a neighbourhood $y \in U \subset M_V$ and identify it with $\mathbb{H}$ such that $y \in M_V$ is mapped into $\mathbf{y} := y^0 + y^1\mathbf{i} + y^2\mathbf{j} + y^3\mathbf{k} \in \mathbb{H}$. Then $B_\varepsilon^4(y) \subset U$ becomes a quaternionic ball $B_\varepsilon^4(\mathbf{y}) \subset \mathbb{H}$ centered about $\mathbf{y}$. Introduce a quaternionic coordinate $\mathbf{x} \in \mathbb{H}$ via $\mathbf{x} := \mathbf{y} + z^1 + z^2\mathbf{j}$ on it. To simplify notation also write g_V for the pullback metric on $B_\varepsilon^4(\mathbf{y})$. There is a bounded positive function a such that $r(\mathbf{x}) = a(\mathbf{x})|\mathbf{x} - \mathbf{y}|$ hence setting $z^1 = x^0 + x^1\mathbf{i}$ and $z^2 = x^2 + x^3\mathbf{i}$ we find in this real coordinate system that

$$\xi^i(\mathbf{x}) = (\delta_j^i + O(|\mathbf{x} - \mathbf{y}|^2)) dx^j \quad \text{and} \quad \xi_i(\mathbf{x}) = (\delta_i^j + O(|\mathbf{x} - \mathbf{y}|^2)) \frac{\partial}{\partial x^j}.$$

Similarly for the harmonic function in (9) we obtain

$$f_{\mathbf{y},\lambda}(\mathbf{x}) = 1 + \frac{\lambda}{|\mathbf{x} - \mathbf{y}|^2} + O(|\mathbf{x} - \mathbf{y}|^{-1}) \quad \text{and} \quad df_{\mathbf{y},\lambda}(\mathbf{x}) = -2\lambda \left(\frac{x^i - y^i}{|\mathbf{x} - \mathbf{y}|^4} \delta_j^i + O(|\mathbf{x} - \mathbf{y}|^{-2}) \right) dx^j.$$

Inserting these expansions into (3) we obtain $a_{\mathbf{y},\lambda} = B_{\mathbf{y},\lambda} + b_{\mathbf{y},\lambda}$ for the perturbation where the singular Euclidean term looks like

$$B_{\mathbf{y},\lambda}(\mathbf{x}) = \text{Im} \frac{\lambda(\bar{\mathbf{x}} - \bar{\mathbf{y}}) d\mathbf{x}}{|\mathbf{x} - \mathbf{y}|^2(\lambda + |\mathbf{x} - \mathbf{y}|^2)}$$

and $b_{\mathbf{y},\lambda}$ is of $O(1)$ hence regular in $B_\varepsilon^4(\mathbf{y})$.

Regarding the spin connection ∇_A , note that the $SU(2)^+$ bundle Σ^+ is trivial over M_V . The metric is anti-self-dual and has vanishing Ricci curvature hence the spin connection is flat and since M_V is simply connected it is just the trivial flat connection. Writing $\nabla_A|_{B_\varepsilon^4(y)} = d + A$ we simply find $A = 0$ in the natural gauge (7).

Putting all of these into (2) we eventually come up in the gauge (7) we use with an expansion of the vector potential as follows:

$$\tilde{A}_{\mathbf{y},\lambda} = B_{\mathbf{y},\lambda} + b_{\mathbf{y},\lambda}$$

whose only singular term is $B_{\mathbf{y},\lambda}$ but still $\tilde{A}_{\mathbf{y},\lambda} \in L_{1,\tilde{A}_{\mathbf{y},\lambda}}^2(\Lambda^1 B_\varepsilon^*(\mathbf{y}) \otimes \text{End}\Sigma^+|_{B_\varepsilon^*(\mathbf{y})})$.

Calculating the curvature we find

$$F_{\tilde{A}_{\mathbf{y},\lambda}} = F_{B_{\mathbf{y},\lambda}} + d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}$$

that is,

$$F_{\tilde{A}_{\mathbf{y},\lambda}}(\mathbf{x}) = \frac{\lambda d\bar{\mathbf{x}} \wedge d\mathbf{x}}{(\lambda + |\mathbf{x} - \mathbf{y}|^2)^2} + d_{B_{\mathbf{y},\lambda}(\mathbf{x})} b_{\mathbf{y},\lambda}(\mathbf{x}) + b_{\mathbf{y},\lambda}(\mathbf{x}) \wedge b_{\mathbf{y},\lambda}(\mathbf{x}) \quad (10)$$

demonstrating that the curvature $F_{B_{\mathbf{y},\lambda}}$ of the Euclidean term as well as $b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}$ are smooth in the ball $B_\varepsilon^4(\mathbf{y})$ meanwhile $d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda}$ is singular. Let $\langle \cdot, \cdot \rangle$ denote the pointwise scalar product and $|\cdot|$ the induced pointwise norm on $\mathfrak{su}(2)^+$ -valued 2-forms associated to the Killing norm on $\mathfrak{su}(2)^+$ and the metric on $\Lambda^2 M_V$. This gives rise to the scalar product $(\cdot, \cdot)_{L^2}$ and norm $\|\cdot\|_{L^2}$ on the punctured ball. Separating the regular and singular terms we obtain

$$\begin{aligned} \|F_{\tilde{A}_{\mathbf{y},\lambda}}\|_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}^2 &= \|F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}\|_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}^2 \\ &+ \|d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda}\|_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}^2 + 2 (F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}, d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda})_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}. \end{aligned}$$

Since the dual of the volume form $*(dV_{g_V}|_{B_\varepsilon(\mathbf{y})}(\mathbf{x}))$ with respect to the flat Hodge star $*$ on $\mathbb{H}$ is $O(|\mathbf{x} - \mathbf{y}|^3)$ we find on the one hand that the regular energy term is of course finite:

$$\|F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}\|_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}^2 = \int_{B_\varepsilon^*(\mathbf{y})} |F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}|^2 dV_{g_V} \leq c_3 \int_0^\varepsilon r^3 dr < +\infty.$$

The singular term $|d_{B_{\mathbf{y},\lambda}(\mathbf{x})} b_{\mathbf{y},\lambda}(\mathbf{x})|$ on the other hand is $O(|\mathbf{x} - \mathbf{y}|^{-1})$ consequently

$$\|d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda}\|_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)}^2 = \int_{B_\varepsilon^*(\mathbf{y})} |d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda}|^2 dV_{g_V} \leq c_4 \int_0^\varepsilon r^{-2} r^3 dr < +\infty.$$

Finally, the cross term also satisfies the estimate (here and only here $|\cdot|$ is the ordinary pointwise absolute value on reals)

$$\begin{aligned} &\left| (F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}, d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda})_{L^2(B_\varepsilon^*(\mathbf{y}), g_V)} \right| \\ &\leq \int_{B_\varepsilon^*(\mathbf{y})} |\langle F_{B_{\mathbf{y},\lambda}} + b_{\mathbf{y},\lambda} \wedge b_{\mathbf{y},\lambda}, d_{B_{\mathbf{y},\lambda}} b_{\mathbf{y},\lambda} \rangle| dV_{g_V} \leq c_5 \int_0^\varepsilon r^{-1} r^3 dr < +\infty. \end{aligned}$$

We conclude that the last two conditions in (5) hold hence the apparent singularity in $\nabla_{\tilde{A}_{y,\lambda}}$ at y can be removed by a gauge transformation via Uhlenbeck's theorem [32] providing us with a smooth anti-self-dual $SU(2)^+$ connection on the whole positive spinor bundle Σ^+ . $\diamond$

Proceeding further we demonstrate that our solutions are anti-instantons, i.e. have finite energy. To carry out this we construct an asymptotic expansion for (9) to see explicitly its fall-off. Viewing asymptotically, a harmonic function with a fixed singularity on (M_V, g_V) is obviously equivalent to pulling all the NUTs in M_V as well as the singular point of the function together; i.e., to construct a harmonic function whose singularity coincides with the singularity of this collapsed space [10]. In light of Lemma 3.1 this corresponds to the case when all the roots of the right hand side in (8) coincide: $p_1 = \dots = p_s = 0$ hence the resulting space $X^* \subset \mathbb{C}^3$ is given by $xy - z^s = 0$. On the other hand, if $(u, v) \in \mathbb{C}^2$ then the polynomials u^s, v^s and uv are invariant under the standard action of $\mathbb{Z}_s \subset SU(2)$ on $\mathbb{C}^2$ and putting $x := u^s, y := v^s, z := uv$ then they also satisfy $xy - z^s = 0$ i.e., X^* is a model in $\mathbb{C}^3$ for the A_{s-1} -singularity $M_V^* := \mathbb{C}^2/\mathbb{Z}_s$.

Removing the origin $\{0\} \in \mathbb{C}^2$ the resulting smooth space $U_V^* := \mathbb{C}^2 \setminus \{0\}/\mathbb{Z}_s$ is topologically a lens space: $U_V^* \cong \mathbb{R}^+ \times L(s, -1)$ where $L(s, -1) \cong S^3/\mathbb{Z}_s$ denotes the usual lens space of $(s, -1)$ -type with its orientation inherited from the complex structure on U_V^* . Regarding U_V^* as a complex line bundle of Chern class $-s$ over S^2 let us denote by (r, τ) polar coordinates on the fibres centered in the singular origin of $\mathbb{C}^2/\mathbb{Z}_s$ and by (Θ, ϕ) the usual spherical coordinates on the base¹. We simply get

$$V(r) = 1 + \frac{s}{2r}, \quad \alpha(\Theta, \phi) = \frac{s}{2} \cos \Theta d\phi$$

consequently the multi-Taub–NUT metric (6) reduces to a singular metric g_V^* on M_V^* whose shape on U_V^* is

$$ds^2 = \frac{2r+s}{2r} (dr^2 + r^2 d\Theta^2 + r^2 \sin^2 \Theta d\phi^2) + \frac{s^2}{4} \frac{2r}{2r+s} (d\tau + \cos \Theta d\phi)^2$$

with parameters

$$0 < r < +\infty, \quad 0 \leq \tau < \frac{4\pi}{s}, \quad 0 \leq \phi < 2\pi, \quad 0 \leq \Theta < \pi. \quad (11)$$

One easily calculates $\det g_V^* = \frac{s^2}{4} (\frac{2r+s}{2})^2 r^2 \sin^2 \Theta$ and by referring to the local expression for the Laplacian of any metric

$$\Delta_g = \sum_{i,j} \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g} g^{ij} \frac{\partial}{\partial x^j} \right),$$

where g^{ij} are the components of the inverse matrix, our singular Laplacian looks like

$$\begin{aligned} \Delta_{g_V^*} = & \frac{2r}{2r+s} \frac{\partial^2}{\partial r^2} + \frac{4}{2r+s} \frac{\partial}{\partial r} + \frac{2(2r+s)^2 \sin^2 \Theta + 2s^2 \cos^2 \Theta}{rs^2(2r+s) \sin^2 \Theta} \frac{\partial^2}{\partial \tau^2} \\ & + \frac{2}{r(2r+s)} \left(\frac{\partial^2}{\partial \Theta^2} + \cot \Theta \frac{\partial}{\partial \Theta} \right) + \frac{2}{r(2r+s) \sin^2 \Theta} \left(\frac{\partial^2}{\partial \phi^2} - 2 \cos \Theta \frac{\partial^2}{\partial \tau \partial \phi} \right). \end{aligned}$$

¹In other words (τ, Θ, ϕ) are Euler coordinates on the lens space and r is the radial coordiante on the cone $\mathbb{C}^2/\mathbb{Z}_s$. For clarity we mention that r defined this way asymptotically accords with the distance function r used in Definition 4.1 while τ here is $\frac{2}{s}$ -times the periodic coordinate τ appeared in (6).

This apparently new Laplacian can be re-expressed in old terms as follows. The lens space with its standard metric induced from the round S^3 has an associated Laplacian

$$\Delta_{L(s,-1)} = \frac{\partial^2}{\partial \Theta^2} + \cot \Theta \frac{\partial}{\partial \Theta} + \frac{1}{\sin^2 \Theta} \left(\frac{\partial^2}{\partial \phi^2} - 2 \cos \Theta \frac{\partial^2}{\partial \tau \partial \phi} + \frac{\partial^2}{\partial \tau^2} \right)$$

in Euler coordinates. By the aid of this we plainly obtain

$$V(r) \Delta_{g_V^*} = \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} + \left(\frac{4}{rs} + \frac{4}{s^2} \right) \frac{\partial^2}{\partial \tau^2} + \frac{1}{r^2} \Delta_{L(s,-1)}.$$

As we have seen, we have to find positive solutions to $\Delta_{g_V^*} f = 0$ or equivalently to

$$V \Delta_{g_V^*} f = 0 \tag{12}$$

with singularity in the origin $r = 0$. These functions can be constructed easily by separating variables. To this end we introduce an orthonormal system of bounded smooth functions on the lens space $L(s, -1)$ by the (three dimensional) spherical harmonics $Y_j^{k,l}$ with $j = 0, 1, \dots$ and $-j \leq k \leq j$, $-\lceil \frac{2j}{s} \rceil \leq l \leq \lceil \frac{2j}{s} \rceil$ (for a fixed j , there are $(2j+1)(\lceil \frac{2j}{s} \rceil + 1)$ independent spherical harmonics on $L(s, -1)$). In Euler coordinates these take the shape

$$Y_j^{k,l}(\tau, \phi, \Theta) = N_j^{k,l,s} e^{i(\frac{ls}{2}\tau + k\phi)} P_j^{k,l}(\cos \Theta)$$

with appropriate normalizing constants $N_j^{k,l,s}$. The generalized associated Legendre polynomial $P_j^{k,l}(x)$ is the unique real analytic solution on $[-1, 1]$ of the following differential equation:

$$\left[(1-x^2) \frac{d^2}{dx^2} - 2x \frac{d}{dx} - \frac{k^2 - x(kls) + \frac{l^2 s^2}{4}}{1-x^2} + j(j+1) \right] P_j^{k,l}(x) = 0.$$

Note that taking $l = 0$ the function $P_j^{k,0}(x)$ reduces to the standard associated Legendre polynomial $P_j^k(x)$ as well as $Y_j^{k,0}(\tau, \phi, \Theta)$ gives back the usual spherical harmonics $Y_j^k(\phi, \Theta)$ on S^2 . The spherical harmonics obey

$$\frac{\partial Y_j^{k,l}}{\partial \tau} = \frac{ils}{2} Y_j^{k,l} \quad \text{and} \quad \Delta_{L(s,-1)} Y_j^{k,l} = -j(j+1) Y_j^{k,l}.$$

In this basis we seek solutions in the form

$$f(r, \tau, \phi, \Theta) = \sum_{j=0}^{\infty} \sum_{k=-j}^j \sum_{l=-\lceil \frac{2j}{s} \rceil}^{\lceil \frac{2j}{s} \rceil} \lambda_j^{k,l} \varrho_j^{k,l}(r) Y_j^{k,l}(\tau, \phi, \Theta)$$

where $\lambda_j^{k,l}$ are complex numbers and $\varrho_j^{k,l}$ are radial functions. Inserting this ansatz into (12) and then exploiting the eigenfunction properties of the spherical harmonics our equation cuts down to a family of ordinary differential equations

$$\left[\frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - \frac{j(j+1)}{r^2} - \frac{sl^2}{r} - l^2 \right] \varrho_j^{k,l}(r) = 0 \tag{13}$$

with $j = 0, 1, \dots$ and $-j \leq k \leq j$, $-\lceil \frac{2j}{s} \rceil \leq l \leq \lceil \frac{2j}{s} \rceil$. The functions $\varrho_j^{k,l}$ are in fact independent of the index k due to the $U(2)$ symmetry of the space (M_V^*, g_V^*) . Two particular solutions are $K_{-j-1}^0(r) = r^{-j-1}$ and $K_j^0(r) = r^j$ if $l = 0$ moreover for $l > 0$ and $r \ll 1$ we know that

$$K_{-j-1}^l(r) = r^{-j-1}(1 + O(r)) \quad \text{and} \quad K_j^l(r) = r^j(1 + O(r))$$

meanwhile for $r \gg 1$ we obtain

$$K_{-j-1}^l(r) = e^{-lr} r^{-\frac{sl}{2}-1}(1 + O(r^{-1})) \quad \text{and} \quad K_j^l(r) = e^{lr} r^{\frac{sl}{2}-1}(1 + O(r^{-1})).$$

Separating variables therefore forces us to put the pointlike singularity of f into the origin of M_V^* showing our approach is consistent. Summing up, the general solution formally looks like

$$f(r, \tau, \phi, \Theta) = \sum_{j=0}^{\infty} \sum_{k=-j}^j \sum_{l=-\lceil \frac{2j}{s} \rceil}^{\lceil \frac{2j}{s} \rceil} \left(\lambda_j^{k,l} K_{-j-1}^l(r) + \mu_j^{k,l} K_j^l(r) \right) Y_j^{k,l}(\tau, \phi, \Theta) \quad (14)$$

with the only singularity in the origin.

Because $\overline{Y_j^{k,l}} = Y_j^{-k,-l}$, a real basis for the solutions is given by

$$\frac{1}{2}(Y_j^{k,l} + Y_j^{-k,-l}) \quad \text{and} \quad \frac{1}{2i}(Y_j^{k,l} - Y_j^{-k,-l}).$$

Moreover from (13) we obtain that $K_{-j-1}^l = K_{-j-1}^{-l}$ and $K_j^l = K_j^{-l}$ consequently if we suppose $\overline{\lambda_j^{k,l}} = \lambda_j^{-k,-l}$ as well as $\overline{\mu_j^{k,l}} = \mu_j^{-k,-l}$ then we come up with formal real solutions in (14).

Taking account that $Y_0^{0,0} = \frac{\sqrt{s}}{4\pi}$ is a constant, the leading $j = 0$ term yields the elementary harmonic function $\lambda r^{-1} + \mu$ (we put simply $\mu_0^{0,0} = \mu$ and $\lambda_0^{0,0} = \lambda$, both real) therefore (9) asymptotically looks like

$$f_{y,\lambda}(x) = 1 + \frac{\lambda}{r} + O\left(\frac{\lambda}{r^{1+\delta}}\right). \quad (15)$$

The leading term is of course positive for $\lambda > 0$. By the aid of this we obtain

Lemma 4.2. *The smooth anti-self-dual connection $\nabla_{\tilde{A}_{y,\lambda}}$ with $y \in M_V$ and $\lambda \in (0, +\infty)$ is subject to both the weak holonomy condition with respect to the trivial flat connection ∇_{Θ} and the rapid decay condition of Definition 2.1 consequently it has not only finite energy but is moreover admissible i.e., its gauge equivalence class gives rise to an admissible anti-instanton on Σ^+ . The energy of this anti-instanton is of one unit.*

Proof. We already have seen that the spin connection on Σ^+ is gauge equivalent to the trivial flat connection ∇_{Θ} .

Consider the neck $W \subset M_V$ of the multi-Taub–NUT manifold and using the gauge (7) along W write $\nabla_{\tilde{A}_{y,\lambda}}|_W = d + \tilde{A}_{y,\lambda}$ with $\tilde{A}_{y,\lambda} = a_{y,\lambda}$ as in (2) where $a_{y,\lambda}$ is the perturbation term involving (9). The asymptotic shape of (9) looks like (15) in light of our previous calculations. This ensures us that if λ is finite and positive, $|d \log f_{\lambda}|_{g_V}$ as well as $|\nabla_{\Theta}(d \log f_{\lambda})|_{g_V}$ are $O(r^{-2})$ along W . Consequently, since

$$|a_{y,\lambda}| = \sqrt{\frac{3}{2}} |d \log f_{y,\lambda}|_{g_V}$$

in the gauge (7), we also obtain that both $|a_{y,\lambda}|$ and $|d_\Theta a_{y,\lambda}|$ are $O(r^{-2})$ along W for all $y \in M_V$ and $0 < \lambda < +\infty$. We see that the first condition in (5) is also satisfied. Inserting this and $F_\Theta = 0$ into (4) we can see that our connection can be viewed as a rapidly decaying perturbation of the trivial connection hence satisfies both the weak holonomy condition with respect to the trivial flat connection as well as the rapid decay condition.

It follows [13, Theorem 4.2] that in this situation the energy of the connection $\nabla_{\tilde{A}_{y,\lambda}}$ is integer hence it must be one because the harmonic function in (9) has only one singularity. $\diamond$

Remark. Before summarizing our findings we may raise the question: what about the higher overtones that is, terms with $j > 0$ in the harmonic expansion (14)? The radial function $r^2 = |u|^2 + |v|^2$ in $\mathbb{C}^2$ is invariant under $\mathbb{Z}_s$ consequently we can talk about lens spaces of radius r centered at the origin of $\mathbb{C}^2/\mathbb{Z}_s$. Take such a lense space with its metric g inherited from the round metric of S^3 . By orthogonality the spherical harmonics obey for $j > 0$ that

$$\frac{1}{2} \int_{L(s,-1)} \left(Y_j^{k,l} + Y_j^{-k,-l} \right) dV_g = \frac{1}{2i} \int_{L(s,-1)} \left(Y_j^{k,l} - Y_j^{-k,-l} \right) dV_g = 0$$

hence these real functions must change sign somewhere along any lense space showing that in fact all individual terms in (14) with $j > 0$ vanish on three dimensional subsets of M_V^* . Additionally we know from the asymptotic formulae for the radial functions K_{-j-1}^l and K_j^l that they behave very differently for small or large r 's consequently the aforementioned three dimensional zero sets of the individual higher overtones cannot be removed by forming particular linear combinations of them. Consequently these harmonic functions represent “non-mild” singularities hence cannot be used to construct anti-instantons.

Let $\nabla_\Gamma|_W$ be a smooth flat $SU(2)^+$ connection on $\Sigma^+|_W$. Then following [13] if $C_0^\infty(\text{End}\Sigma^+)$ is the space of smooth, compactly supported endomorphisms of the positive chiral spin bundle then define the L^2 gauge group $\mathcal{G}_{\Sigma^+}$ as the $L_{2,\Gamma}^2$ completion of the space

$$\{\gamma - 1 \in C_0^\infty(\text{End}\Sigma^+) \mid \|\gamma - 1\|_{L_{2,\Gamma}^2(M_V, g_V)} < +\infty, \gamma \in C^\infty(\text{Aut}\Sigma^+) \text{ a.e.}\}$$

Let us denote by $\mathcal{M}(e, \Gamma)$ the *framed moduli space* consisting of pairs $([\nabla_A], \Gamma)$, the $\mathcal{G}_{\Sigma^+}$ gauge equivalence classes of smooth irreducible $SU(2)^+$ anti-self-dual connections on Σ^+ of energy $e < +\infty$, decaying rapidly and obeying the weak holonomy condition with respect to $\nabla_\Gamma|_W$ and a fixed gauge Γ at infinity preserved by $\mathcal{G}_{\Sigma^+}$. Associated to this space one has the *unframed moduli space* $\widehat{\mathcal{M}}(e, \Gamma)$ consisting of the $\mathcal{G}_{\Sigma^+}$ equivalence classes $[\nabla_A]$ only. In other words it is formed by dividing via the group

$$\widehat{\mathcal{G}}_{\Sigma^+} \cong \mathcal{G}_{\Sigma^+} \times SU(2)^+ \quad (16)$$

of gauge transformations tending to any fixed element $\gamma_0 \in SU(2)^+$ (not to the identity only) at infinity.

In particular we can consider the spaces $\mathcal{M}(k, \Theta)$ and $\widehat{\mathcal{M}}(k, \Theta)$ corresponding to the trivial flat connection ∇_Θ ; in this case the energy is an integer $k \in \mathbb{N}$, cf. [13, Theorem 4.2]. It was demonstrated in [13, Theorem 3.2] that these spaces are smooth manifolds and [13, Theorem 4.2] ensures us that $\dim \mathcal{M}(k, \Theta) = 8k$ and therefore $\dim \widehat{\mathcal{M}}(k, \Theta) = 8k - 3$.

Putting together our findings sofar as well as noting that all of our anti-instantons are irreducible [12, Theorem 5.1] we obtain the following

Theorem 4.1. *Let (M_V, g_V) be the multi-Taub–NUT space with $s > 0$ NUTs equipped with an orientation coming from any of its complex structures in the hyper-Kähler family.*

Then there exists a 5 real parameter family of unframed, non-gauge equivalent, irreducible, smooth $SU(2)^+$ anti-instantons

$$\{[\nabla_{\tilde{A}_{y,\lambda}}] \mid (y, \lambda) \in M_V \times (0, +\infty)\}$$

on the positive chiral spinor bundle Σ^+ which satisfy the weak holonomy condition with respect to the trivial flat connection and also satisfy the rapid decay condition. All these anti-instantons have unital energy and provide us with an open subset of (one connected component of) the unframed moduli space $\widehat{\mathcal{M}}(1, \Theta)$. $\diamond$

Theorem 4.1 can be regarded as a familiar “collar theorem” for the 1-anti-instanton moduli space over the multi-Taub–NUT space. The patient reader may write down these solutions explicitly by inserting the explicit Green functions [30, Equation (18)] of Page into (9) and then by the aid of the gauge (7) construct the corresponding connection 1-form via (2). However the quite complicated result is not informative.

Rather we turn attention to a puzzling feature of the description provided by Theorem 4.1. This is the limit $\lambda \rightarrow +\infty$. In this case (9) up to reparametrization remains meaningful and provides us with the positive minimal Green function $G(\cdot, y)$. It readily follows from the proofs of Lemmata 4.1 and 4.2 that the resulting anti-self-dual connection $\nabla_{\tilde{A}_{y,+\infty}}$ gives rise to an unframed anti-instanton $[\nabla_{\tilde{A}_{y,+\infty}}]$ on Σ^+ obeying the weak holonomy condition with respect to ∇_Θ and the rapid decay condition hence these solutions are expected to complete the moduli space.

Indeed, for a fixed $y \in M_V$ taking the limit $\lambda \rightarrow +\infty$ (especially in (10)) the proof of Lemma 4.1 continues to hold hence the corresponding anti-self-dual connection is smooth everywhere on Σ^+ .

Additionally, in light of (15) the asymptotics of $G(\cdot, y)$ looks like $1/r$ therefore the proof of Lemma 4.2 also can be repeated if we notice that this time $|a_y|$ and $|d_\Theta a_y|$ are $O(r^{-1})$ only. Hence $a_y \notin L^2_{\frac{1}{2},1,\Theta}(\Lambda^1 M \otimes \text{End} \Sigma^+)$ however it follows from a direct calculation that the corresponding curvature still belongs to the weighted Sobolev space: $F_{\tilde{A}_y} \in L^2_{\frac{1}{2}}(\Lambda^2 M \otimes \text{End} \Sigma^+)$. We conclude that $[\nabla_{\tilde{A}_{y,+\infty}}]$ provides us with an admissible unframed anti-instanton with unit energy on the positive spin bundle; more precisely it decays rapidly and one can also check that the weak holonomy condition with respect to the same trivial flat connection ∇_Θ is fulfilled. If $y \in M_V$ is not a NUT then the corresponding anti-instanton is irreducible otherwise reducible to $U(1)$ and these Abelian solutions span the full L^2 cohomology of the multi-Taub–NUT geometry as it was stated in [12, Theorem 5.1].

Therefore if $y \in M_V$ is not a NUT then these limiting irreducible solutions are expected to give the completion of the collar $M_V \times (0, +\infty)$ to the full unframed moduli space in the unconcentrated or “centerless” regime—or at least one connected component of the moduli space emerges this way. Since it is natural to include the reducible solutions corresponding to the NUTs into $\widehat{\mathcal{M}}(1, \Theta)$ as well, for simplicity we shall continue to denote (one connected component of) this extended moduli space by $\widehat{\mathcal{M}}(1, \Theta)$ from now on and note that it is not a manifold anymore: usual conical singularities appear around the reducible points as we know from the general theory but will also see shortly.

We also conjecturally note that in our opinion not only a connected component but the whole moduli space of unital energy admissible anti-instantons with trivial holonomy arises this way.

Theorem 4.2. *Consider the multi-Taub–NUT space (M_V, g_V) with $s > 0$ NUTs $p_1, \dots, p_s \in M_V$, equipped with the natural orientation as above. Then for (one connected component of) the unframed moduli space of unital energy $SU(2)^+$ admissible anti-instantons decaying rapidly to the trivial flat connection on Σ^+ we find*

$$\widehat{\mathcal{M}}(1, \Theta) \cong (M_V \times (0, +\infty]) / \sim \quad (17)$$

where the equivalence relation $\sim$ means that $M_V \times \{+\infty\}$ is pinched into $\mathbb{R}^3$ by collapsing the S^1 -isometry orbits of (M_V, g_V) .

Consequently there exists a singular fibration

$$\Phi : \widehat{\mathcal{M}}(1, \Theta) \longrightarrow \mathbb{R}^3 \quad (18)$$

with generic fibers homeomorphic to the open 2-ball B^2 and as many as s singular fibers homeomorphic to the semi-open 1-ball $(0, +\infty]$. Therefore (one connected component of) the moduli space is contractible and in particular is orientable.

The images of the points $(p_i, +\infty)$ in $\widehat{\mathcal{M}}(1, \Theta)$ with $i = 1, 2, \dots, s$ represent reducible anti-instantons and $\widehat{\mathcal{M}}(1, \Theta)$ around these points looks like a cone over $\mathbb{CP}^2$ or $\overline{\mathbb{CP}}^2$ (depending on the orientation).

Proof. In Sect. 5 we will prove that if $y \in M_V$ and $e^{i\tau}y$ is its image by an element of the S^1 isometry group of (M_V, g_V) then the corresponding two anti-instantons with $(y, \lambda = +\infty)$ and $(e^{i\tau}y, \lambda = +\infty)$ are actually gauge equivalent unital energy, rapidly decaying anti-instantons with trivial holonomy in infinity. On the other hand we know from [13, Theorem 3.2] that away from the reducible points the moduli space is a finite dimensional, possibly non-connected manifold without boundary.

Consequently we are forced to look at our moduli space as follows. Fix a point $y \in M_V$ different from any NUT and consider the S^1 isometry orbit through y : it is parameterized by the cyclic coordinate $\tau \in [0, 2\pi)$ of the local coordinate system on (M_V, g_V) introduced in Sect. 3; as well as take the concentration parameter $\lambda \in (0, +\infty)$. Then by Theorem 4.1, associated to any $y \in M_V$, we obtain a subset of the moduli space parameterized by (τ, λ) . We regard this as a punctured open B^2 such that the points with $\lambda = 0$ are supposed to represent the boundary while $\lambda = +\infty$ the center of this (τ, λ) -ball. In this picture we interpret the existence of smooth solutions with $\lambda = +\infty$ as filling in the punctured (τ, λ) -ball over $y \in M_V$ with the anti-instanton $[\nabla_{\tilde{A}_{y, +\infty}}]$ corresponding to the limit $\lambda \rightarrow +\infty$. In other words, this anti-instanton represents the center of the unpunctured (τ, λ) -ball that consequently will be homeomorphic to an open B^2 over any $y \in M_V$ different from the NUTs.

Since the NUTs $p_1, \dots, p_s \in M_V$ are the only fixed points of the isometry group action on (M_V, g_V) it follows that over these points the moduli space is parameterized by $\lambda \in (0, +\infty]$ only hence is a semi-open B^1 . In this case the “center” is the endpoint $\lambda = +\infty$.

Therefore completing our collar $M_V \times (0, +\infty)$ with these limiting solutions, the moduli space takes the shape as in (17) where the equivalence relation $\sim$ means that $M_V \times \{+\infty\}$ is collapsed into $\mathbb{R}^3$ by shrinking all the isometry orbits of (M_V, g_V) into single points respectively and admits a singular fibration (18) whose generic fiber is an open B^2 and with s pieces of singular fibers homeomorphic to a semi-open B^1 . In particular $\widehat{\mathcal{M}}(1, \Theta)$ is contractible hence orientable.

We already know that the only reducible solutions are the images $r_i \in \widehat{\mathcal{M}}(1, \Theta)$ of the points $(p_i, +\infty)$ with $i = 1, 2, \dots, s$. We will study the structure of the moduli space around these points. Take a NUT $p_i \in M_V$ and consider $B^5(r_i) \subset \widehat{\mathcal{M}}(1, \Theta)$. We will construct a decomposition

$\partial\overline{B}^5(r_i) = X_1 \cup X_2$ of the boundary as follows. It is clear that if y is not a NUT and the image of $(y, +\infty)$ in $\widehat{\mathcal{M}}(1, \Theta)$ is m_i then $B^5(m_i) \cong B^5$, an open 5-ball. Let us decompose its closure as $\overline{B}^3 \times \overline{B}^2$ where B^2 is the aforementioned (τ, λ) -disk and B^3 is a ball around the image of y in $\mathbb{R}^3$. Then $\partial\overline{B}^5 \cong (S^2 \times \overline{B}^2) \cup (\overline{B}^3 \times S^1)$ which is of course an S^4 glued together from $S^2 \times \overline{B}^2$ and $\overline{B}^3 \times S^1$ along their common boundary $S^2 \times S^1$. Now let us move y toward a particular NUT p_i ; then one of the $\overline{B}^2$'s in the $S^2 \times \overline{B}^2$ component of the above decomposition, namely that one which is moved exactly into the position of the NUT, gets shrink to a $\overline{B}^1$ in a manner such that meanwhile its radius (measured by λ) is kept constant, its circumference (measured by τ) gets vanish. Put

$$X_1 \cong S^2 \times \overline{B}^2 / \sim_1$$

corresponding to this collapsed space. We obviously find

$$H^j(X_1; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } j = 0, 2 \\ 0 & \text{otherwise.} \end{cases}$$

Since this movement also collapses an S^1 , the boundary in the other component $\overline{B}^3 \times S^1$ of the previously collapsed $\overline{B}^2$, we also get a corresponding collapsed space

$$X_2 \cong \overline{B}^3 \times S^1 / \sim_2 .$$

Since the generator of $\pi_1(\overline{B}^3 \times S^1) \cong \mathbb{Z}$ is just killed out in X_2 , we also obtain

$$H^j(X_2; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } j = 0 \\ 0 & \text{otherwise.} \end{cases}$$

It is also clear that $X_1 \cap X_2$ is nothing else than a 3-sphere with exactly two antipodal points pinched consequently

$$H^j(X_1 \cap X_2; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } j = 0, 3 \\ 0 & \text{otherwise.} \end{cases}$$

Recalling the Mayer–Vietoris sequence of the decomposition $\partial\overline{B}^5(r_i) = X_1 \cup X_2$ which looks like

$$\cdots \rightarrow H^j(\partial\overline{B}^5(r_i); \mathbb{Z}) \rightarrow H^j(X_1; \mathbb{Z}) \oplus H^j(X_2; \mathbb{Z}) \rightarrow H^j(X_1 \cap X_2; \mathbb{Z}) \rightarrow H^{j+1}(\partial\overline{B}^5(r_i); \mathbb{Z}) \rightarrow \cdots$$

we end up with

$$H^j(\partial\overline{B}^5(r_i); \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } j = 0, 2, 4 \\ 0 & \text{otherwise.} \end{cases}$$

However from this we obtain via Freedman's classification [14] that $\partial\overline{B}^5(r_i) \cong \mathbb{C}P^2$ or $\overline{\mathbb{C}P}^2$ depending on the orientation put onto $\widehat{\mathcal{M}}(1, \Theta)$. Therefore the unframed moduli space about a reducible point looks like a cone over one of these spaces as stated. $\diamond$

We make a comment on the higher-energy moduli spaces including anti-instantons with trivial holonomy in infinity. It is obvious that instead of the functions (9) we can use harmonic functions with a finite number of isolated singularities yielding

Theorem 4.3. *The moduli spaces $\mathcal{M}(k, \Theta)$ hence $\widehat{\mathcal{M}}(k, \Theta)$ are non-empty for all $k \in \mathbb{N}$. $\diamond$*

For sake of completeness without proof we exhibit here the degenerate $s = 0$ case as well i.e., the flat $\mathbb{R}^3 \times S^1$ whose associated instantons also referred to as *calorons*. This space is flat hence conformal rescaling works. Using coordinates $(x^1, x^2, x^3, \tau) = (x, \tau)$ on $\mathbb{R}^3 \times S^1$ we have an obvious five parameter family of positive harmonic functions (9) with Green functions in a rather explicit form [30]

$$G((x, \tau_x), (y, \tau_y)) = \frac{1}{16\pi^2|x-y|} \frac{\tanh|x-y|}{1 - \frac{\cos(\tau_x - \tau_y)}{\cosh|x-y|}} = \frac{1}{4\pi^2} \sum_{k=-\infty}^{+\infty} \frac{1}{|x-y|^2 + (\tau_x - \tau_y + 2\pi k)^2}.$$

Making use of [13, Theorem 2.3] to describe the holonomy issue in this situation as well as [13, Theorem 4.1] which provides us with the dimension moreover taking into account that there are no reducible $SU(2)^\pm$ instantons on this space we state

Theorem 4.4. *Take the flat $\mathbb{R}^3 \times S^1$ with an arbitrary fixed orientation. Then any non-trivial $SU(2)^\pm$ instanton on $\Sigma^\pm$ is irreducible moreover the rapidly decaying ones obey the weak holonomy condition with respect to some flat connection ∇_Γ on the neck and their energies are always non-negative integers $k \in \mathbb{N}$. The corresponding framed moduli spaces, if not empty, have dimensions $\dim \mathcal{M}(\Gamma, k) = 8k$ while $\dim \widehat{\mathcal{M}}(k, \Gamma) = 8k - 3$ for the unframed ones.*

Assume the asymptotic flat connection is the trivial one ∇_Θ and let $\mathcal{M}(1, \Theta)$ be the corresponding framed moduli space. Then for (one connected component of) the unframed space we find

$$\widehat{\mathcal{M}}(1, \Theta) \cong (\mathbb{R}^3 \times S^1 \times (0, +\infty]) / \sim \cong \mathbb{R}^3 \times B^2$$

moreover the higher energy framed moduli spaces $\mathcal{M}(k, \Theta)$ hence $\widehat{\mathcal{M}}(k, \Theta)$ are also non-empty with $k \in \mathbb{N}$ arbitrary. $\diamond$

5 Completion of the moduli space

In this closing section our aim is to prove that conformal rescaling gives rise to at least one connected component of unital energy, rapidly decaying anti-instantons obeying the weak holonomy condition with respect to the trivial flat connection (however we note again that in our opinion this moduli space is connected).

Our starting point is the construction of the twistor space of the multi-Taub–NUT geometry by the aid of Hitchin [22] and then a certain compactification of it. This enables us to identify the harmonic functions used so far with elements of certain Ext groups on the compactified twistor space following Atiyah [2]. These Ext groups also provide us with twisted holomorphic vector bundles over the original non-compact twistor space and the corresponding untwisted ones represent our anti-instantons, in the spirit of the Atiyah–Ward correspondence, in a new form.

To get these twisted vector bundles more explicitly we investigate that real structure on the non-compact twistor space which leads to the multi-Taub–NUT geometry and also construct the corresponding real twistor lines; our twisted vector bundles then, referring to Serre’s method, can be constructed via sections vanishing along these real lines. The transition functions of these twisted bundles corresponding to pure Green functions, i.e., in the $\lambda \rightarrow +\infty$ limit can be constructed rather explicitly. We will find that the twisted bundles corresponding to $G(\cdot, y)$ and $G(\cdot, e^{i\tau}y)$ are actually isomorphic hence the corresponding anti-instantons must be gauge equivalent verifying our picture on the moduli space presented in Sect. 4.

To begin with, we claim that

Lemma 5.1. *The twistor space Z of any multi-Taub–NUT space (M_V, g_V) with s NUTs admits a smooth complex compactification $\overline{Z}$. This compactification arises by adding $s + 3$ pieces of various Hirzebruch surfaces to Z in a suitable way.*

Proof. Fix a generic configuration of s points $q_1, \dots, q_s \in \mathbb{R}^3$ and consider the corresponding multi-Taub–NUT space (M_V, g_V) . Pick a direction e_1 in $\mathbb{R}^3$ not parallel with any straight line segment ℓ_{ij} connecting q_i and q_j and put the corresponding (generic) integrable complex structure J_{e_1} onto M_V from the hyper-Kähler family. We know that (M_V, J_{e_1}) is biholomorphic to the algebraic surface $X \subset \mathbb{C}^3$ given by (8). However there are at most $2\binom{s}{2} = s(s-1)$ exceptional directions to choose for e_1 such that the corresponding complex manifolds are *not* biholomorphic to X .

Introducing homogeneous coordinates $([x : u], [y : v], [z : w]) \in \mathbb{C}P_x^1 \times \mathbb{C}P_y^1 \times \mathbb{C}P_z^1$ equation (8) can be made homogeneous of degrees $(1, 1, s)$ respectively as follows:

$$xyw^s - uv \prod_{i=1}^s (z - wp_i) = 0. \quad (19)$$

Let us denote the resulting compact complex surface in $\mathbb{C}P_x^1 \times \mathbb{C}P_y^1 \times \mathbb{C}P_z^1$ by $\overline{X}^*$. We claim that $\overline{X}^*$ is smooth except one point. Indeed, let $0_x := [0 : 1]$ and $\infty_x := [1 : 0]$ denote the origin and the infinity of $\mathbb{C}P_x^1$ respectively and in the same fashion introduce the notation $0_y, \infty_y \in \mathbb{C}P_y^1$ and $0_z, \infty_z \in \mathbb{C}P_z^1$. Then in the compactification $\mathbb{C}^3 \subset \mathbb{C}P_x^1 \times \mathbb{C}P_y^1 \times \mathbb{C}P_z^1$ infinity is represented by the bouquet

$$\infty := (\{\infty_x\} \times \mathbb{C}P_y^1 \times \mathbb{C}P_z^1) \vee (\mathbb{C}P_x^1 \times \{\infty_y\} \times \mathbb{C}P_z^1) \vee (\mathbb{C}P_x^1 \times \mathbb{C}P_y^1 \times \{\infty_z\})$$

and one obtains that $\overline{X}^* \cap \infty$ consists of four lines, that is $\overline{X}^* = X \sqcup (\ell_1 \cup \ell_2 \cup \ell_3 \cup \ell_4)$ with

$$\begin{aligned} \ell_1 &:= \{\infty_x\} \times \{0_y\} \times \mathbb{C}P_z^1, & \ell_2 &:= \{\infty_x\} \times \mathbb{C}P_y^1 \times \{\infty_z\}, \\ \ell_3 &:= \mathbb{C}P_x^1 \times \{\infty_y\} \times \{\infty_z\}, & \ell_4 &:= \{0_x\} \times \{\infty_y\} \times \mathbb{C}P_z^1. \end{aligned}$$

These lines are not disjoint, they intersect each other as follows:

$$\ell_1 \cap \ell_2 = (\infty_x, 0_y, \infty_z), \quad \ell_2 \cap \ell_3 = (\infty_x, \infty_y, \infty_z), \quad \ell_3 \cap \ell_4 = (0_x, \infty_y, \infty_z).$$

By calculating the gradient of the left hand side in (19) we see that the only point where $\overline{X}^*$ is singular is the intersection $\ell_2 \cap \ell_3$ where $\overline{X}^*$ locally looks like

$$uv - \prod_{i=1}^s w(1 - wp_i)^{-1} = 0 \quad (20)$$

i.e., it possesses a classical A_{s-1} quotient singularity.

(M_V, g_V) being simply connected with vanishing Ricci curvature, its twistor space Z admits a holomorphic fibration over $\mathbb{C}P^1$ consequently we would like to regard equation (19) as a set of equations parameterized by a projective line. Therefore we proceed as follows (cf. [22]). Consider the holomorphic fiber bundle

$$\overline{\pi} : P(H^k \oplus H^0) \oplus P(H^l \oplus H^0) \oplus P(H^m \oplus H^0) \longrightarrow \mathbb{C}P^1 \quad (21)$$

whose fibers are $F_x \times F_y \times F_z$ with $F_i \cong \mathbb{C}P^1$ ($i = x, y, z$) as before. Take the subbundle $\bar{\pi}_x : P(H^k \oplus H^0) \rightarrow \mathbb{C}P^1$ with $F_x \cong \mathbb{C}P^1$. Referring to the isomorphism

$$P(H^0 \oplus H^{-k}) \xrightarrow{\otimes H^k} P(H^k \oplus H^0)$$

take the canonical section $\infty_x := (1, 0)$ of $P(H^0 \oplus H^{-k})$ and regard its image as a divisor which we also denote by ∞_x ; one then has an associated line bundle $L_x := [\infty_x]$ and corresponding sheaf $\mathcal{O}_{\text{rel},x}(1)$ on $P(H^0 \oplus H^{-k})$. This line bundle has a canonical section $u \in \mathcal{O}_{\text{rel},x}(1)$ satisfying $(u) = \infty_x$. We also have a canonical section $0_x := (0, 1)$ of $P(H^k \oplus H^0)$ which gives rise to the bundle $L_x \otimes \bar{\pi}_x^* H^k$ and a canonical section x which is an element of $\mathcal{O}_{\text{rel},x}(1) \otimes \mathcal{O}_{P(H^k \oplus H^0)}(k)$ where $\mathcal{O}_{P(H^k \oplus H^0)}(k)$ is the associated sheaf of $\bar{\pi}_x^* H^k$. Take the fibration

$$\bar{\pi}_{y,z} : P(H^k \oplus H^0) \oplus P(H^l \oplus H^0) \oplus P(H^m \oplus H^0) \longrightarrow P(H^k \oplus H^0)$$

whose fibers are $F_y \times F_z$. Pulling back x, u we therefore obtain elements

$$\bar{\pi}_{y,z}^* x \in \bar{\pi}_{y,z}^* \mathcal{O}_{\text{rel},x}(1) \otimes \bar{\pi}_{y,z}^* \mathcal{O}_{P(H^k \oplus H^0)}(k), \quad \bar{\pi}_{y,z}^* u \in \bar{\pi}_{y,z}^* \mathcal{O}_{\text{rel},x}(1).$$

In the same fashion we construct

$$\bar{\pi}_{x,z}^* y \in \bar{\pi}_{x,z}^* \mathcal{O}_{\text{rel},y}(1) \otimes \bar{\pi}_{x,z}^* \mathcal{O}_{P(H^l \oplus H^0)}(l), \quad \bar{\pi}_{x,z}^* v \in \bar{\pi}_{x,z}^* \mathcal{O}_{\text{rel},y}(1)$$

and finally

$$\bar{\pi}_{x,y}^* z \in \bar{\pi}_{x,y}^* \mathcal{O}_{\text{rel},z}(1) \otimes \bar{\pi}_{x,y}^* \mathcal{O}_{P(H^m \oplus H^0)}(m), \quad \bar{\pi}_{x,y}^* w \in \bar{\pi}_{x,y}^* \mathcal{O}_{\text{rel},z}(1).$$

For simplicity we shall denote them as x, u, y, v, z, w respectively and observe that they provide us with coordinates on the total space of the fibration (21).

The situation gets simplified somewhat by virtue of the following observation. It is not difficult to check that the bundle $\bar{\pi}_{y,z}^* (\bar{\pi}_x^* H^k)$ on the total space of (21) depends only on k and not on the way we pulled it back from $\mathbb{C}P^1$. Consequently

$$\bar{\pi}_{y,z}^* (\bar{\pi}_x^* H^k) \cong \bar{\pi}_{x,y}^* (\bar{\pi}_z^* H^k) \cong \bar{\pi}_{x,z}^* (\bar{\pi}_y^* H^k) \cong \bar{\pi}^* H^k$$

and the same is true for the other bundles H^l and H^m . Write

$$W := P(H^k \oplus H^0) \oplus P(H^l \oplus H^0) \oplus P(H^m \oplus H^0)$$

for the total space of (21) and put

$$L^{a_0, a_1, a_2, a_3} := \bar{\pi}^* H^{a_0} \otimes \bar{\pi}_{y,z}^* L_x^{a_1} \otimes \bar{\pi}_{x,z}^* L_y^{a_2} \otimes \bar{\pi}_{x,y}^* L_z^{a_3}$$

with $a_i \in \mathbb{Z}$; we will also use the convenient notation for the corresponding sheaf:

$$\mathcal{O}_W(a_0, a_1, a_2, a_3) := \mathcal{O}_W(a_0) \otimes \bar{\pi}_{y,z}^* \mathcal{O}_{\text{rel},x}(a_1) \otimes \bar{\pi}_{x,z}^* \mathcal{O}_{\text{rel},y}(a_2) \otimes \bar{\pi}_{x,y}^* \mathcal{O}_{\text{rel},z}(a_3)$$

(therefore $\mathcal{O}_W(a_0) = \mathcal{O}_W(a_0, 0, 0, 0)$). Then $x \in \mathcal{O}_W(k, 1, 0, 0)$ and $u \in \mathcal{O}_W(0, 1, 0, 0)$, etc. and if we regard $p_i \in \mathcal{O}_W(m, 0, 0, 0)$ we obtain

$$xyw^s \in \mathcal{O}_W(k+l, 1, 1, s) \quad \text{and} \quad uv \prod_{i=1}^s (z - wp_i) \in \mathcal{O}_W(sm, 1, 1, s)$$

consequently setting $k + l = sm$ we end up with a well-defined map $\overline{P} : W \longrightarrow L^{sm,1,1,s}$ where formally $\overline{P}$ is given by the left hand side of (19) and a singular model for the compactified twistor space $\overline{Z}^* \subset W$ is provided by the hypersurface $\overline{P} = 0$ hence restricting (21) it admits a holomorphic fibration:

$$\pi : \overline{Z}^* \longrightarrow \mathbb{CP}^1. \quad (22)$$

Note that $\overline{Z}^*$ has an obvious singularity along the line ℓ which is the section passing through the singular points $(\infty_x, \infty_y, \infty_z)_{[a:b]} \in \overline{X}_{[a:b]}^* = \pi^{-1}([a : b])$ for an $[a : b] \in \mathbb{CP}^1$. However $\overline{Z}^*$ has more singularities, namely the points $(0, 0, p_i([a : b]))$ where exactly two roots coincide: $p_i([a : b]) = p_j([a : b])$. (We may assume that the configuration is generic hence no more than two roots coincide at the same time.) Hence these points represent rational double points in $\overline{Z}^*$. Therefore the number of such bad points is $m \binom{s}{2} = m \cdot \frac{1}{2}s(s-1)$ since $p_i \in \mathcal{O}_W(m, 0, 0, 0)$. The geometric origin of these bad points is clear: they just correspond to the special complex structures analyzed in Lemma 3.1 under which (M_V, J_{e_1}) is *not* biholomorphic to (8). Since the number of these complex structures is $s(s-1)$ as we have seen, we have to set $m = 2$ yielding $k + l = 2s$ and since x and y play symmetric roles, we take $k = l = s$ in (21).

We finally desingularize $\overline{Z}^*$ to obtain a smooth compact complex 3-manifold $\overline{Z}$ whose finite part $Z \subset \overline{Z}$ provides us with a non-singular model for the twistor space of (M_V, g_V) . In light of Lemma 3.1 we simply do this by replacing the fibers $\overline{X}_{[a:b]}^*$ of (22) with their non-singular blowups for all $[a : b] \in \mathbb{CP}^1$. Let us denote the resulting smooth fiber by $\overline{X}_{[a:b]}$. Clearly, for a generic value of $[a : b]$ we have to just remove the A_{s-1} -singularity at infinity from $\overline{X}_{[a:b]}^*$ therefore we have to attach $4 + s - 1 = s + 3$ lines in an appropriate way:

$$\overline{X}_{[a:b]} = X_{[a:b]} \sqcup (\ell_1 \cup \ell_2 \cup \ell'_1 \cup \dots \cup \ell'_{s-1} \cup \ell_3 \cup \ell_4) \quad (23)$$

otherwise we have to carry out an additional blowup in the point $(0, 0, p_j(a, b)) \in \overline{X}_{[a:b]}^*$ as well. Since these fibers are diffeomorphic to each other, we end up with a smooth holomorphic fibration that for simplicity we continue to denote as

$$\pi : \overline{Z} \longrightarrow \mathbb{CP}^1. \quad (24)$$

We therefore conclude that the compactified smooth $\overline{Z}$ arises by adding $s + 3$ Hirzebruch surfaces to the smooth twistor space Z of (M_V, g_V) as follows:

$$\overline{Z} = Z \sqcup (S_1 \cup S_2 \cup S'_1 \cup \dots \cup S'_{s-1} \cup S_3 \cup S_4).$$

Two of them, S_1 and S_4 stemming from the lines ℓ_1 and ℓ_4 in $\overline{X}^*$, are biholomorphic to $P(H^2 \oplus H^0)$ while S_2 and S_3 given by ℓ_2 and ℓ_3 are biholomorphic to $P(H^s \oplus H^0)$. To determine the type of the exceptional ones, notice that the local resolution of the singularity (20) is given by defining the new variables $u_s := w^{-s}u, v_s := v, w_s := w$, whence the equation of the proper transform becomes

$$u_s v_s - \prod_{i=1}^s (1 - w_s p_i)^{-1} = 0.$$

This means that we add a family of projective lines parametrized by $\mathbb{CP}^1$ with sections w_s . As w_s is untwisted in the base direction, this implies that the Hirzebruch surfaces $S'_1, \dots, S'_{s-1}$ we add through this process are trivial $\mathbb{CP}^1 \times \mathbb{CP}^1$'s. $\diamond$

Remark. It is worth comparing the complex compactification (23) with the simple smooth real compactification of M_V in [11, 13].

Lemma 5.2. *The middle sheaf cohomology groups of the compactified smooth twistor space $\overline{Z}$ satisfy*

$$H^1(\overline{Z}; \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C} \quad \text{and} \quad H^2(\overline{Z}; \mathcal{O}_{\overline{Z}}(-2)) \cong 0.$$

Proof. First notice that referring to (21) for all $[a : b] \in \mathbb{CP}^1$ the fiber $\overline{X}_{[a:b]}^* := \pi^{-1}([a : b])$ is a rational surface. Indeed, consider a general $[a : b] \in \mathbb{CP}^1$, where the fiber is smooth except at infinity. Then $\overline{X}_{[a:b]}^* \cong \overline{X}^*$ is given by equation (19) for distinct values of $p_j = p_j([a : b])$. Consider the map

$$\overline{X}^* \longrightarrow \mathbb{CP}_x^1 \times \mathbb{CP}_z^1, \quad ([x : u], [y : v], [z : w]) \longmapsto ([x : u], [z : w]).$$

Over a given $([x : u], [z : w]) \in \mathbb{CP}_x^1 \times \mathbb{CP}_z^1$ there exists a unique $[y : v] \in \mathbb{CP}_y^1$ such that $([x : u], [y : v], [z : w]) \in \overline{X}^*$, except in the case $xw^s = u \prod (z - wp_j) = 0$ when any $[y : v]$ is a solution. It readily follows that these exceptional points are precisely

$$\{(0_x, [p_1 : 1]), \dots, (0_x, [p_s : 1]), (\infty_x, \infty_z)\} \in \mathbb{CP}_x^1 \times \mathbb{CP}_z^1. \quad (25)$$

Furthermore, all the points of $\overline{X}^*$ are non-singular except the isolated hence codimension 2 surface singularity of type A_{s-1} at $(\infty_x, \infty_y, \infty_z) \in \mathbb{CP}_x^1 \times \mathbb{CP}_y^1 \times \mathbb{CP}_z^1$ which is mapped onto (∞_x, ∞_z) in $\mathbb{CP}_x^1 \times \mathbb{CP}_z^1$. It follows that $\overline{X}_{[a:b]}^*$ is isomorphic to the blowup of $\mathbb{CP}_x^1 \times \mathbb{CP}_z^1$ in the points (25) outside of a codimension 2 subset. The same argument works for the special set of $[a : b] \in \mathbb{CP}^1$ where $\overline{X}_{[a:b]}^*$ admits a further rational double point, if we add this singular point to the codimension 2 subset above. Therefore $\overline{X}_{[a:b]}^*$ is rational. Since by Lemma 5.1 for all $[a : b] \in \mathbb{CP}^1$ the non-singular fiber $\overline{X}_{[a:b]}^*$ of (24) arises by blowing further up $\overline{X}_{[a:b]}^*$ in finitely many points, we conclude that $\overline{X}_{[a:b]}$ is also rational.

Next, notice that $\mathcal{O}_{\overline{Z}}(-2) = \pi^* \mathcal{O}(-2)$ is trivial along the fibers of (24). Rationality implies in particular that for all $[a : b]$ and $1 \leq q \leq 2$ one has

$$H^q \left(\overline{X}_{[a:b]}; (\pi|_{\overline{X}_{[a:b]}})^* \mathcal{O}(-2) \right) \cong H^q \left(\overline{X}_{[a:b]}; \mathcal{O}_{\overline{X}_{[a:b]}} \right) = 0,$$

so $R^q \pi_* \pi^* \mathcal{O}(-2)$ is the zero sheaf. On the other hand $R^0 \pi_* \pi^* \mathcal{O}(-2) \cong \mathcal{O}(-2)$ by the projection formula consequently

$$H^0(\mathbb{CP}^1; R^0 \pi_* \pi^* \mathcal{O}(-2)) = 0 \quad \text{and} \quad H^1(\mathbb{CP}^1; R^0 \pi_* \pi^* \mathcal{O}(-2)) = \mathbb{C}.$$

The second level of the Leray spectral sequence associated to (24) and the sheaf $\pi^* \mathcal{O}(-2)$ is given by $E_2^{p,q} = H^p(\mathbb{CP}^1; R^q \pi_* \pi^* \mathcal{O}(-2))$. Taking into account our calculations so far, the only non-trivial part is a one-dimensional space in bidegree $(1, 0)$:

$$\begin{array}{c|cc} & 0 & 0 \\ q & 0 & 0 \\ & 0 & \mathbb{C} \\ \hline & p & \end{array}$$

implying that the spectral sequence collapses at this step and the statement follows. $\diamond$

Keeping these results in mind now we recall the notion of the “global Ext” groups $\text{Ext}^k(V; \mathcal{R}, \mathcal{S})$ over a complex manifold V with two coherent sheaves $\mathcal{R}$ and $\mathcal{S}$ on it. The background material can be found in [16]. These objects are appropriate generalizations of the sheaf cohomology groups over complex manifolds. For instance if $\mathcal{R} = \mathcal{O}_V$ then we know that

$$\text{Ext}^k(V; \mathcal{O}_V, \mathcal{S}) \cong H^k(V; \mathcal{S}). \quad (26)$$

If V is compact n dimensional then there is a generalization of the Serre duality theorem for coherent sheaves, called the *Grothendieck duality theorem*:

$$H^k(V; \mathcal{S}) \cong (\text{Ext}^{n-k}(V; \mathcal{S}, \mathcal{K}_V))^* \quad (27)$$

and in the same fashion if $\mathcal{S} = \mathcal{O}(E)$ is locally free then the *Serre duality theorem* implies that

$$(\text{Ext}^{n-k}(V; \mathcal{O}(E), \mathcal{K}_V))^* \cong H^{n-k}(V; \mathcal{O}(E^*) \otimes \mathcal{K}_V). \quad (28)$$

We will be particularly interested in $\text{Ext}^1(V; \mathcal{R}, \mathcal{S})$ which classifies extensions of coherent sheaves over V :

$$0 \longrightarrow \mathcal{S} \longrightarrow \mathcal{E} \longrightarrow \mathcal{R} \longrightarrow 0.$$

Following Atiyah [2] we obtain:

Lemma 5.3. *Let $\mathcal{J}_Y$ denote the ideal sheaf of holomorphic functions on $\overline{Z}$ vanishing on a fixed real line $Y \subset Z \subset \overline{Z}$. Then*

$$\text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C}^2.$$

Proof. Note that $\mathcal{O}_Y(-2) \cong \mathcal{K}_Y$, the canonical sheaf of Y . Let $j : Y \rightarrow \overline{Z}$ be the inclusion and denote by $j_*\mathcal{K}_Y$ the extension of $\mathcal{K}_Y$ by zero over $\overline{Z}$. Hence $j_*\mathcal{K}_Y$ is a coherent sheaf on $\overline{Z}$. Applying (27) on $\overline{Z}$ in the last step we obtain

$$H^k(Y; \mathcal{K}_Y) \cong H^k(\overline{Z}; j_*\mathcal{K}_Y) \cong (\text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{K}_Y, \mathcal{K}_{\overline{Z}}))^*$$

and additionally we know that

$$\text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{K}_Y, \mathcal{K}_{\overline{Z}}) \cong \text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{O}_Y \otimes \overline{\pi}^*\mathcal{K}_Y, \mathcal{K}_{\overline{Z}}) \cong \text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{O}_Y, \overline{\pi}^*\mathcal{K}_Y^* \otimes \mathcal{K}_{\overline{Z}}).$$

We already know that $(\overline{\pi}^*\mathcal{K}_Y^* \otimes \mathcal{K}_{\overline{Z}})|_Y$ is non-canonically isomorphic to H^{-2} , therefore $\mathcal{O}_{\overline{Z}}(-2)$ is an extension of $(\overline{\pi}^*\mathcal{K}_Y^* \otimes \mathcal{K}_{\overline{Z}})|_Y$ over the whole $\overline{Z}$ that is,

$$\text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{O}_Y, \overline{\pi}^*\mathcal{K}_Y^* \otimes \mathcal{K}_{\overline{Z}}) \cong \text{Ext}^{3-k}(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)).$$

Therefore, taking into account that $H^2(Y; \mathcal{K}_Y) \cong 0$ and $H^1(Y; \mathcal{K}_Y) \cong \mathbb{C}$ we conclude that

$$\text{Ext}^1(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \cong 0 \quad \text{and} \quad \text{Ext}^2(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C}.$$

Referring to (26) we know $\text{Ext}^i(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \cong H^i(\overline{Z}; \mathcal{O}_{\overline{Z}}(-2))$ consequently Lemma 5.2 additionally yields

$$\text{Ext}^1(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C} \quad \text{and} \quad \text{Ext}^2(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \cong 0.$$

Now consider the short exact sequence of coherent sheaves over $\overline{Z}$:

$$0 \longrightarrow \mathcal{J}_Y \longrightarrow \mathcal{O}_{\overline{Z}} \longrightarrow j_*\mathcal{O}_Y \longrightarrow 0.$$

A segment of the associated long exact sequence of the global Ext groups with $\mathcal{O}_{\overline{Z}}(-2)$ looks like

$$\begin{aligned} \dots &\longrightarrow \text{Ext}^1(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \\ &\longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \xrightarrow{\delta} \text{Ext}^2(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \\ &\longrightarrow \text{Ext}^2(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \dots \end{aligned}$$

which gives

$$0 \longrightarrow \mathbb{C} \longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \mathbb{C} \longrightarrow 0 \quad (29)$$

providing the result. $\diamond$

After these algebro-geometric preliminaries the time has come to relate our considerations so far with anti-instantons over a multi-Taub–NUT space.

First we establish a link with harmonic functions. The embedding $i : Z \setminus Y \subset \overline{Z}$ together with (26) implies a homomorphism

$$i^* : \text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \text{Ext}^1(Z \setminus Y; \mathcal{J}_Y, \mathcal{O}_{Z \setminus Y}(-2)) \cong H^1(Z \setminus Y, \mathcal{O}_{Z \setminus Y}(-2)).$$

However for a fixed real line $Y \subset Z$ representing the point $y \in M_V$ the group $H^1(Z \setminus Y; \mathcal{O}_{Z \setminus Y}(-2))$ gives rise to an isomorphism

$$T : H^1(Z \setminus Y; \mathcal{O}_{Z \setminus Y}(-2)) \cong \text{Ker } \Delta_{g_V}|_{M_V \setminus \{y\}}$$

which is the Penrose transform in this case. Consequently one can think of the elements of $\text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2))$ as harmonic functions. Therefore Lemma 5.3 provides us with a distinguished 2-parameter family of complex-valued harmonic functions on $M_V \setminus \{y\}$. These functions can easily be found by observing that the positive minimal Green function with singularity in $y \in M_V$ of Definition 4.1 provides us with a splitting of $\text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C}^2$ as a vector space into two one dimensional summands. More precisely (29) gives

$$0 \longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \xrightarrow{\delta} \text{Ext}^2(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow 0$$

and the Green function defines a map

$$\text{Ext}^2(\overline{Z}; j_*\mathcal{O}_Y, \mathcal{O}_{\overline{Z}}(-2)) \longrightarrow \text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)),$$

so that all elements of the middle term canonically can be written in the form $\alpha + \beta$ and the first factor α comes from $\text{Ext}^1(\overline{Z}; \mathcal{O}_{\overline{Z}}, \mathcal{O}_{\overline{Z}}(-2))$ which is independent of Y hence corresponds to a *constant* harmonic function on (M_V, g_V) .

Consequently we obtain the following

Theorem 5.1. (Atiyah, 1981) *Fix a point $y \in M_V$ of the multi-Taub–NUT space (M_V, g_V) and consider the associated real line $Y \subset Z$ and the group $\text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2)) \cong \mathbb{C}^2$. Then any element of this group can canonically be written in the form $\mu + \lambda G(\cdot, y)$ where $G(\cdot, y)$ is the unique minimal positive Green function concentrated at $y \in M_V$ and $\mu, \lambda \in \mathbb{C}$ are constants. $\diamond$*

Remark. Notice that in fact the above theorem holds for any anti-self-dual non-parabolic manifold whose twistor space admits a compactification satisfying Lemma 5.2.

One can then see that the positive harmonic functions of (9)—from which our anti-instantons $[\nabla_{\tilde{A}_{y,\lambda}}]$ stem—can also be regarded as elements of $\text{Ext}^1(\overline{Z}; \mathcal{J}_Y, \mathcal{O}_{\overline{Z}}(-2))$ hence they describe extensions of two coherent sheaves

$$0 \longrightarrow \mathcal{O}_{\overline{Z}}(-2) \longrightarrow \overline{\mathcal{F}}(-1) \longrightarrow \mathcal{J}_Y \longrightarrow 0.$$

Restricting this to Z , the resulting sheaves $\mathcal{F}(-1)$ will be locally free hence provide us with rank 2 holomorphic vector bundles $F(-1)$ over Z .

Our aim is now to understand those twisted vector bundles together with their canonical sections among the aforementioned vector bundles over Z which are associated to pure Green functions. We want to show that for two points $y, e^{i\tau}y \in M_V$ on the same S^1 -orbit, these twisted vector bundles are isomorphic.

For this purpose, it is sufficient to work over the open twistor space $Z \subset \overline{Z}$, because we are only interested in the real lines lying over points $y \in M_V$ at finite distance. Therefore we take a closer look at this space.

Restricting our construction in Lemma 5.1 and in particular the singular fibration (22) to the finite part we obtain a model for the non-compact singular twistor space $\pi : Z^* \rightarrow \mathbb{CP}^1$ as the hypersurface $P = 0$ given by

$$P : H^s \oplus H^s \oplus H^2 \rightarrow H^{2s}, \quad P(x, y, z) = xy - \prod_{j=1}^s (z - p_j). \quad (30)$$

The choice $k = l = s$ and $m = 2$ in Lemma 5.1 is also dictated by the requirement that the normal bundle of a real line $Y \subset Z$ must be isomorphic to $H \oplus H$ and Z must possess a non-trivial real structure.

Regarding the former property, to describe the normal bundle of a real line $Y \subset Z$ is equivalent to describe that of a generic real line $Y \subset Z^*$. Since it is a section of the bundle $\pi : Z^* \rightarrow \mathbb{CP}^1$ its normal bundle is the restriction of the tangent bundle of $H^s \oplus H^s \oplus H^2$ to a generic (non-singular) line i.e., the kernel of the map $(P_x, P_y, P_z) : T(H^s \oplus H^s \oplus H^2) \rightarrow TH^{2s}$. This shows that $\langle c_1(N_Y), [Y] \rangle = s + s + 2 - 2s = 2$ moreover the type of the normal bundle is stable under small perturbations yielding $N_Y \cong H \oplus H$. An important consequence is that since $K_Z|_Y \cong K_Y \otimes \Lambda^2 N_Y^*$ we therefore find $K_Z \cong \pi^* H^{-4}$ for the canonical bundle of Z as we already noted in Sect. 2. Notice that this relation does not extend to the compactification $\overline{Z}$. Indeed, for every real line Y at finite distance the relation $N_Y \cong H \oplus H$ continues to hold, but because the fibers of the map (24) become compact, in $K_{\overline{Z}}$ there will be a twist coming from the canonical bundle of $\overline{X}$ as well. This explains the asymmetry between the middle cohomology groups in Lemma 5.2.

Next we move to the construction of the real structure on Z following [10] — originally due to [22, 24] — which gives rise to the multi-Taub–NUT geometry. The idea is to twist the natural real structure on Z which gives rise to the A_{s-1} ALE geometries [22] with sections of a certain line bundle L^c which lives on the total space of the line bundle

$$\pi_z : H^2 \rightarrow \mathbb{CP}^1.$$

It is again enough to work with the singular model (30) however now we will think of the fibration $\pi : H^s \oplus H^s \oplus H^2 \rightarrow \mathbb{CP}^1$ as $\pi : H^s \oplus H^s \rightarrow H^2$.

For the definition of L^c , let the base space $\mathbb{CP}^1$ of Z^* be covered by the affine open sets U_a and U_b with coordinates $[a : b]$ satisfying $a \neq 0$ and $b \neq 0$ respectively and write $U'_a = \pi_z^{-1}(U_a)$ and $U'_b = \pi_z^{-1}(U_b)$ for the open subsets of H^2 covering the affine subsets of $\mathbb{CP}^1$ moreover denote

by z any point in the total space of H^2 . Therefore we obtain coordinate systems $([1 : \frac{b}{a}], z)$ and $([\frac{a}{b} : 1], z)$ on U'_a and U'_b respectively. Define the line bundle L^c with a complex parameter c by the transition function

$$g : U'_a \cap U'_b \longrightarrow \mathbb{C}^*, \quad g(z, [a : b]) := e^{-cz \frac{b}{a}}.$$

It is clear that L^0 is just the trivial bundle over H^2 .

Note that $H^2 \cong T\mathbb{C}P^1$ can be naturally identified with the space of oriented affine lines in $\mathbb{R}^3$ (with a fixed origin) as follows: if $\ell \subset \mathbb{R}^3$ is an oriented affine line and $\ell(t) \in \ell$ is its point, $t \in \mathbb{R}$, then clearly exist unique vectors $\mathbf{u}, \mathbf{v}$ with $|\mathbf{u}| = 1$ and $\mathbf{u} \cdot \mathbf{v} = 0$ such that $\ell(t) = \mathbf{v} + t\mathbf{u}$. Regarding $\mathbf{u}$ as a point on S^2 and $\mathbf{v}$ a vector in its tangent space we obtain the isomorphism. There is a natural real structure on the space of oriented real lines namely $(\mathbf{u}, \mathbf{v}) \mapsto (-\mathbf{u}, \mathbf{v})$ which gives rise to the antipodal map on $\mathbb{C}P^1$ inside H^2 as a zero section consequently we also obtain a real structure on the space of holomorphic sections $r : H^0(\mathbb{C}P^1; \mathcal{O}(2)) \rightarrow H^0(\mathbb{C}P^1; \mathcal{O}(2))$. In the same fashion we obtain real structures $r : H^0(H^2; \mathcal{O}(H^s \otimes L^c)) \rightarrow H^0(H^2; \mathcal{O}(H^s \otimes L^{-c}))$ satisfying $r^2 = (-1)^s \text{Id}$. A section is called *real* if it is invariant under this map. Recall that the real structure on Z is induced by the antipodal map on $\mathbb{C}P^1$. Therefore the real structure on Z^* is defined to be

$$\tau_c(x, y, z) := ((-1)^s r(y), r(x), -r(z)). \quad (31)$$

We find the corresponding real lines by a factorization method [22, 24]. If $z \in H^2$ then there is a unique oriented affine line $\ell_z \subset \mathbb{R}^3$ corresponding to it with a distinguished point $\ell(0) \in \mathbb{R}^3$ (depending on the choice of origin in $\mathbb{R}^3$). This point can be characterized by the set of all lines in $\mathbb{R}^3$ intersecting it which provides us with a real section in $H^0(\mathbb{C}P^1; \mathcal{O}(2)) \cong \mathbb{C}^3$ hence is of the form $z([a : b]) = \alpha a^2 + 2\beta ab - \bar{\alpha} b^2$ with $\alpha \in \mathbb{C}$ and $\beta \in \mathbb{R}$. Therefore we have two ways of looking at the same object: we can regard z either as a point on the total space $z \in H^2$ or as a real section $\alpha a^2 + 2\beta ab - \bar{\alpha} b^2 \in H^0(\mathbb{C}P^1; \mathcal{O}(2))$. Of course, quite tautologically speaking, the section z passes through the point z corresponding to it.

Regarding z as a real section $z([a : b]) = \alpha a^2 + 2\beta ab - \bar{\alpha} b^2$ now, this homogeneous polynomial is canonically indentified with the single-valued polynomial in the affine coordinate $\frac{a}{b}$ over U_b :

$$z|_{U_b} \left(\frac{a}{b} \right) = \alpha \left(\frac{a}{b} \right)^2 + 2\beta \left(\frac{a}{b} \right) - \bar{\alpha}.$$

Now we define a nowhere vanishing section η^c of L^c by

$$\eta^c(z, [a : b]) := \begin{cases} e^{c(\alpha(\frac{a}{b}) + \beta)} & \text{on } U'_b; \\ g(z, [a : b]) e^{c(\alpha(\frac{a}{b}) + \beta)} & \text{on } U'_a. \end{cases}$$

It is well-defined since it looks like $\exp c(\alpha(\frac{a}{b}) + \beta)$ on U'_b i.e., where $b \neq 0$ and looks like $\exp c(\bar{\alpha}(\frac{b}{a}) - \beta)$ on U'_a i.e., where $a \neq 0$.

If $p_j([\frac{a}{b} : 1])|_{U_b} = \alpha_j(\frac{a}{b}) + 2\beta_j(\frac{a}{b}) - \bar{\alpha}_j$ and $\zeta([\frac{a}{b} : 1])|_{U_b} = \alpha(\frac{a}{b}) + 2\beta(\frac{a}{b}) - \bar{\alpha}$ then the roots $\rho_j, \sigma_j \in \mathbb{C} \subset \mathbb{C}P^1$ of $\zeta - p_j = 0$ look like

$$\rho_j := \frac{-(\beta - \beta_j) - \sqrt{(\beta - \beta_j)^2 + |\alpha - \alpha_j|^2}}{\alpha - \alpha_j}, \quad \sigma_j := \frac{-(\beta - \beta_j) + \sqrt{(\beta - \beta_j)^2 + |\alpha - \alpha_j|^2}}{\alpha - \alpha_j}$$

and the real lines appear by simply factorizing in a τ_c -invariant way the equation (30) and imposing the reality condition (this already has been done for ζ). They are of the form

$$\begin{cases} \xi(\zeta, [a : b]) = A\eta^c(\zeta, [a : b]) \prod_{j=1}^s (a - \rho_j b) \\ v(\zeta, [a : b]) = B\eta^{-c}(\zeta, [a : b]) \prod_{j=1}^s (a - \sigma_j b) \\ \zeta([a : b]) = \alpha a^2 + 2\beta ab - \bar{\alpha} b^2. \end{cases}$$

The coefficients A, B here are arbitrary constants satisfying $AB = \prod(\alpha - \alpha_i)$, the leading coefficient of $\prod(\zeta - p_i)$. As this latter number is non-zero for $\alpha \neq \alpha_j$, say B can be expressed in terms of A and α . But the reality constraint moreover implies

$$|A|^2 = \prod_j \left((\beta - \beta_j) + \sqrt{(\beta - \beta_j)^2 + |\alpha - \alpha_j|^2} \right).$$

Hence the set of these real lines is locally parameterized² by $\arg A \in S^1$ and $(\operatorname{Re} \alpha, \operatorname{Im} \alpha, \beta) \in \mathbb{R}^3$ and provides us with a local chart of M_V as it was constructed in Sect. 3. The choice $c = 0$ gives rise to the ALE A_{s-1} geometries while $c > 0$ real provides us with the ALF A_{s-1} i.e., the multi-Taub–NUT spaces. Hence from now on we will restrict to $c = 1$ in accord with previous sections.

The next step is to express the transition matrices for the vector bundle $F_{\alpha, \beta, A}(-1)$ on Z as well as its section $s_{\alpha, \beta, A} \in H^0(Z; \mathcal{O}(F_{\alpha, \beta, A}(-1)))$, associated to the canonical element

$$G(\cdot, y_{\alpha, \beta, A}) \in \operatorname{Ext}^1(Z \setminus Y_{\alpha, \beta, A}; \mathcal{I}_{Y_{\alpha, \beta, A}}, \mathcal{O}_{Z \setminus Y_{\alpha, \beta, A}}(-2))$$

representing the Green function. Here, we follow again [2] and first work over Z^* . Set

$$\begin{aligned} f &:= x - \xi \\ g &:= y - v \\ h &:= z - \zeta. \end{aligned}$$

By definition, the real twistor line $Y_{\alpha, \beta, A}$ is given by the equations

$$f = g = h = 0.$$

Define open subsets U_ξ and U_v by $\xi \neq 0$ and $v \neq 0$ respectively. Then, restricted to $U_\xi \cap Z^*$ the real line is given by the complete intersection $f = h = 0$ as well as restricted to $U_v \cap Z^*$ it is the complete intersection $g = h = 0$. Furthermore, for all generic ζ the polynomials $\zeta - p_i$ and $\zeta - p_j$ do not have common roots for $i \neq j$, so U_ξ and U_v cover Z^* . We restrict to such choices α, β, A , but by continuity our results continue to hold in the general case too. Let us now set

$$\theta := \frac{xy - \xi v}{h},$$

a polynomial in a, b . Straightforward verification yields the identities

$$f = -\frac{x}{v}g + \frac{\theta}{v}h \quad \text{and} \quad h = -\frac{h}{v}g + \frac{y}{v}h.$$

²Note that the way of assigning the roots of $\zeta - p_j$ to ξ and v is well-defined only up to an action of the Galois group of the problem which explains that we have a local parameterization only.

Obviously f, g and h are well-defined sections, vanishing on $Y_{\alpha,\beta,A}$, of $H^s \otimes L^1$, $H^s \otimes L^{-1}$ and H^2 respectively.

Hence, we can define a rank 2 vector bundle $F_{\alpha,\beta,A}(-1)$ on Z^* by gluing the sections f and h of the bundle $(H^s \otimes L^1) \oplus H^2$ restricted to U_ξ with the sections g and h of the bundle $(H^s \otimes L^{-1}) \oplus H^2$ on U_v using the gluing matrix

$$M_{\alpha,\beta,A} := \frac{1}{v} \begin{pmatrix} -x & \theta \\ -h & y \end{pmatrix} \quad (32)$$

on $U_\xi \cap U_v$. Moreover, we get a section $s_{\alpha,\beta,A}$ of this vector bundle by setting it to be equal to (f, h) on U_ξ and to (g, h) on U_v . This section vanishes precisely on $Y_{\alpha,\beta,A}$.

Recall also that S^1 acts on M_V as identified with the space of real sections by multiplication on the component A , or equivalently in view of the identity $AB = \prod(\alpha - \alpha_i)$, by inverse multiplication on B . Our aim is to show that if we consider the twistor line associated to a point $(\alpha, \beta, e^{i\tau}A)$, then the vector bundle $F(-1)$ constructed for this point will be isomorphic to the one corresponding to (α, β, A) . But the gluing matrix (32) for these former data is related to that of the latter ones via

$$M_{\alpha,\beta,e^{i\tau}A} = e^{-i\tau} M_{\alpha,\beta,A}.$$

It follows that the change of trivializations $(f, h) \mapsto (e^{i\tau}f, e^{i\tau}h)$ on U_ξ induces an isomorphism between the bundles $F_{\alpha,\beta,A}(-1)$ and $F_{\alpha,\beta,e^{i\tau}A}(-1)$.

As we have seen we obtain Z from the singular space Z^* by blowing it up in finitely many points. Carrying out this procedure and then pulling back the bundles just constructed we come up with isomorphic vector bundles $F_{\alpha,\beta,A}(-1)$ and $F_{\alpha,\beta,e^{i\tau}A}(-1)$ over the regular (and non-compact) twistor space Z .

Consider the untwisted vector bundle $F_{\alpha,\beta,A} = F_{\alpha,\beta,A}(-1) \otimes H$. First of all, it is clear that it is a holomorphic rank 2 vector bundle over Z .

Secondly, the restriction of $F_{\alpha,\beta,A}$ to any real line in Z is holomorphically trivial. This follows since the bundle we have constructed just corresponds to the Green function as in [2] and this function is everywhere positive as we already noted [17].

Thirdly, the anti-linear map (31) induces a real structure on $F_{\alpha,\beta,A}(-1)$ over τ_1 simply because τ_1 switches $H^s \otimes L^1$ and $H^s \otimes L^{-1}$ while maps H^2 to H^2 . Therefore, since over $\mathbb{CP}^1$ the bundle H carries a symplectic structure [18], the untwisted bundle $F_{\alpha,\beta,A} = F_{\alpha,\beta,A}(-1) \otimes H$ also carries a symplectic structure $\tilde{\tau}_1 : F_{\alpha,\beta,A} \rightarrow F_{\alpha,\beta,A}$ lying over τ_1 .

Since the same conclusions are also true for $F_{\alpha,\beta,e^{i\tau}A}$ it then follows that the corresponding untwisted vector bundles satisfy the three conditions of the Atiyah–Ward correspondence (cf. the summary of the Atiyah–Ward correspondence in Sect. 2) hence they indeed give rise to anti-instantons. Since $F_{\alpha,\beta,A}$ and $F_{\alpha,\beta,e^{i\tau}A}$ are isomorphic, we therefore conclude that the limiting anti-instantons are gauge equivalent i.e., $[\nabla_{\tilde{A}_{y,+\infty}}] = [\nabla_{\tilde{A}_{e^{i\tau}y,+\infty}}]$ as desired.

References

- [1] Anderson, M.T., Kronheimer, P.B., LeBrun, C.: *Complete Ricci-flat manifolds with infinite topological type*, Comm. Math. Phys. **125**, 637-642 (1989);
- [2] Atiyah, M.F.: *Green's functions for self-dual four-manifolds*, Mathematical analysis and applications, Part A, Advances in mathematics supplementary studies **7A**, 129-158 (1981);

- [3] Atiyah, M.F., Hitchin, N.J.: *The geometry and dynamics of magnetic monopoles*, Princeton University Press, Princeton (1988);
- [4] Atiyah, M.F., Hitchin, N.J., Singer, I.M.: *Self-duality in four-dimensional Riemannian geometry*, Proc. Roy. Soc. London **A362**, 425–461 (1978);
- [5] Atiyah, M.F., Ward, R.: *Instantons and algebraic geometry*, Comm. Math. Phys. **55**, 117–124 (1977);
- [6] Besse, A.L.: *Einstein Manifolds*, Springer-Verlag, Berlin (1987);
- [7] Biquard, O., Jardim, M.: *Asymptotic behaviour and moduli space of double periodic instantons*, Journ. Eur. Math. Soc. **3**, 335–375 (2001);
- [8] Cherkis, S.: *Moduli spaces of instantons on the Taub–NUT space*, preprint, 25 pp., arXiv: [hep-th/0805.1245](#) (2008);
- [9] Cherkis, S., Hitchin, N.J.: *Gravitational instantons of type D_k* , Comm. Math. Phys. **260**, 299–317 (2005);
- [10] Etesi, G.: *Classification of 't Hooft instantons over multi-centered gravitational instantons*, Nucl. Phys. **B662**, 511–530 (2003);
- [11] Etesi, G.: *The topology of asymptotically locally flat gravitational instantons*, Phys. Lett. **B641**, 461–465 (2006);
- [12] Etesi, G., Hausel, T.: *On Yang–Mills-instantons over multi-centered gravitational instantons*, Comm. Math. Phys. **235**, 275–288 (2003), arXiv: [hep-th/0207196](#);
- [13] Etesi, G., Jardim, M.: *Moduli spaces of self-dual connections over asymptotically locally flat gravitainal instantons*, Comm. Math. Phys. **280**, 285–313 (2008), arXiv: [math.DG/0608597](#);
- [14] Freedman, M.H.: *The topology of four-manifolds*, Journ. Diff. Geom. **17**, 357–454 (1982);
- [15] Gibbons, G.W., Hawking, S.W.: *Gravitational multi-instantons*, Phys. Lett. **B78**, 430–432 (1978);
- [16] Griffiths, Ph., Harris, J.: *Principles of Algebraic Geometry*, John Wiley & Sons, Inc., New York (1984);
- [17] Grigor'yan, A.A.: *The existence of positive fundamental solutions of the Laplace equation on Riemannian manifolds* (Russian), Math. Sb. (S.N.) **128 (170)**, 354–363, 446 (1985);
- [18] Hartshorne, R.: *Stable vector bundles and instantons*, Comm. Math. Phys. **59**, 1–15 (1978);
- [19] Hausel, T., Hunsicker, E., Mazzeo, R.: *Hodge cohomology of gravitational instantons*, Duke Math. Journ. **122**, 485–548 (2004);
- [20] Hawking, S.W.: *Gravitational instantons*, Phys. Lett. **A60**, 81–83 (1977);
- [21] Hebey, E.: *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*, Courant Lecture Notes **5**, AMS (2000);

- [22] Hitchin, N.J.: *Polygons and gravitons*, Math. Proc. Camb. Phyl. Soc. **85**, 465-476 (1979);
- [23] Hitchin, N.J.: *Linear field equations on self-dual spaces*, Proc. Roy. Soc. London **A370**, 173-191 (1980);
- [24] Hitchin, N.J.: *Twistor construction of Einstein metrics*, in: Global Riemannian geometry (Durham, 1983), 115-125, Ellis Horwood Ser.: Math. Appl., Horwood, Chichester (1984);
- [25] Jackiw, R., Nohl, C., Rebbi, C.: *Conformal properties of pseudo-particle configurations*, Phys. Rev. **D15**, 1642-1646 (1977);
- [26] Kronheimer, P.B.: *The construction of ALE spaces as hyper-Kähler quotients*, Journ. Diff. Geom. **29** 665-683 (1989);
- [27] Kronheimer, P.B., Nakajima, H.: *Yang–Mills instantons on ALE gravitational instantons*, Math. Ann. **288**, no.2., 263-607 (1990);
- [28] LeBrun, C.: *Complete Ricci-flat Kähler metrics on $\mathbb{C}^n$ need not be flat*, Proc. Symp. Pure Math. **52** (2), 297-304 (1991);
- [29] Minerbe, V.: *On some asymptotically flat manifolds with non-maximal volume growth*, preprint, 49 pp., arXiv: math.DG/0709.1084 (2007);
- [30] Page, D.N.: *Green’s function for gravitational multi-instantons*, Phys. Lett. **B85**, 369-372 (1979);
- [31] Penrose, R.: *Nonlinear gravitons and curved twistor theory*, Gen. Rel. Grav. **7**, 31-52 (1976);
- [32] Uhlenbeck, K.K.: *Removable singularities in Yang–Mills fields*, Comm. Math. Phys. **83**, 11-29 (1982);
- [33] Varopoulos, N.: *Potential theory and diffusion on Riemannian manifolds*, Conference on harmonic analysis in honor of Antoni Zygmund (Chicago, Ill. 1981), Vol. I, II, 821-837, Wadsworth Math. Ser., Wadsworth, Belmont, USA (1983).